CUMULATIVE AUTHOR INDEX
Environmental Entomology
Volume 39, 2010

A

Abel, C. A., 1085, 1111
Adams, A. S., 406
Adams, S. M., 406
Adler, L. S., 1953
Aguilar, A., 827
Ahmad, A. H., 210
Akhtar, Z. R., 715
Aldrich, J. R., 610
Ali, A., 210
Aliakbarpour, H., 1409
Allahyari, H., 1586
Allan, S. A., 1608
Allemand, R., 535
Allen, C., 1085
Almada, E. D., 1134
Al-Shami, S., 210
Aluko, G. K., 1715
Al-Wahaibi, A. K., 15
Amézquita, S., 328
Amarasekare, K. G., 1506
Anderson, R. S., 1848
Anderson, S. L., 190
Anduaga, S., 1165
Angelella, G. M., 505
Araujo, E. L., 1457
Arbogast, R. T., 1731
Aukema, B. H., 874, 898

B

Bae, Y. J., 278
Bahlai, C. A., 50
Bai, Y. Y., 243
Baker, T. C., 169
Bakker, J. D., 1159
Balciunas, J., 114, 841
Barber, N. A., 492
Barraclough, E. I., 1359
Barrett, J. P., 633
Barro, P., 1652
Basoalto, E., 1399
Bauce, E., 498
Bauer, L. S., 1513
Bauman, T. A., 633
Bean, D. W., 1666
Beers, E. H., 286
Belovsky, G. E., 312
Bencivenga, N., 1170
Bennett, K. C., 1631
Bento, F.d.M.M., 1315
Bento, J.M.S., 678
Berenbaum, M.R., 1291, 1922
Berkebile, D. R., 1101
Bethke, J. A., 159
Beuzelin, J. M., 528
Bi, M.-J., 883
Biddinger, D. J., 2045
Björklund, N., 898
Blair, D., 1122
Blake, A. J., 1195
Blankenship, E. E., 202
Bledsoe, L., 484
Boetel, M. A., 1545
Bosch, J., 149
Botha, A.-M., 1206
Branson, D. H., 346, 1441
Breidenbaugh, M. S., 591
Brewer, M. J., 31
Brightwell, R.j., 1659
Broadbent, A. B., 1570
Broce, A. B., 1101
Brock, C., 1961
Brousseau, P.-M., 1151
Brown, G. C., 1309, 1788
Brown, J. K., 1028
Brown, M. D., 545
Brown, M. W., 134
Buczkowski, G., 1936
Bueno, V.H.P., 1500
Burgess, E.P.J., 1359
Butler, R. C., 303
Byun, B.-K., 337

C

Cafaro, M. J., 105
Calatayud, P.-A., 57
Campbell, J. B., 1101
Campbell, J. F., 919
Campos, M., 79
Candia, O., 1276
Cao, W.-J., 232
Carneiro, M.A.A., 849, 1134
Carney, V. A., 738, 1666
Carpaneto, G. M., 1756
Carpenter, J. E., 1884
Carreira, V. P., 865
Carver, S., 798
Cavanagh, A. F., 1953
Chaŕez-Moreno, C. K., 1178
Chabi-Olaye, A., 468
Chacón, E., 1848
Chapman, R. B., 303
Chen, H.-S., 1021
Chen, M., 715, 1369
Chen, P., 1378
Chen, W., 1705
Chen, Wenlong., 1545
Chen, X. X., 1369
Chen, X.-X., 2039
Chen, Y., 715, 1369
Cheng, D.-F., 1737
Cheng, J. A., 243
Cheng, R.-X., 389
Cherry, R. H., 1012, 2025
Chinnasamy, D., 1980
Chiozza, M. V., 856
Choate, B., 1428, 1447
Chu, D., 1028
Chun, J. H., 337
Clayton, M. K., 874
Cloutier, C., 1151
Cockfield, S. D., 286
Cokendolpher, J. C., 1428
Collins, J. A., 1428
Colunga-Garcia, M., 31
Cook, S. P., 821
Coop, L. B., 2006
Cooper, W. R., 223, 787
Corcuera, P., 320
Corio, C., 865
Corn, J. G., 164
Cornelissen, T., 849
Cory, J. S., 459
Costa, C.M.Q., 359
Costa, M.l.Z., 1315
Costa-Lima, T. C., 195
Cotes, B., 79
Cottrell, T. E., 610
Coyle, D. R., 1242, 1539
Currie, C. R., 105, 406
Curtis, P. S., 1810

D

Daane, K. M., 1232, 1264
Dale, P., 88
Dalin, P., 1666
Daniels, J. C., 1961
Danne, A., 970
Davidson, L. N., 576
Davis, A.L.V., 811
Davis, G. A., 440
Debano, S. J., 1
Defoliart, L. S., 1643
Delatte, J., 1715
Delpuech, J. M., 535
Delucia, E. H., 1291
Deschodt, C. M., 811
De M.M. Bento, F., 1315
De La Mora, A., 1473
De Vries, R. S. M., 459
De Szalay, F. A., 591
De Lyra, M. C. C. P., 1178
Dickey, A. M., 1037
Dieng, H., 1409
Dillwith, J. W., 223
Dively, G. P., 707
Donaldson, J., 1561
Dorn, S., 561

CUMULATIVE AUTHOR INDEX
Environmental Entomology
Volume 39, 2010

A

Abel, C. A., 1085, 1111
Adams, A. S., 406
Adams, S. M., 406
Adler, L. S., 1953
Aguilar, A., 827
Ahmad, A. H., 210
Akhtar, Z. R., 715
Aldrich, J. R., 610
Ali, A., 210
Aliakbarpour, H., 1409
Allahyari, H., 1586
Allan, S. A., 1608
Allemand, R., 535
Allen, C., 1085
Almada, E. D., 1134
Al-Shami, S., 210
Aluko, G. K., 1715
Al-Wahaibi, A. K., 15
Amézquita, S., 328
Amarasekare, K. G., 1506
Anderson, R. S., 1848
Anderson, S. L., 190
Anduaga, S., 1165
Angelella, G. M., 505
Araujo, E. L., 1457
Arbogast, R. T., 1731
Aukema, B. H., 874, 898

B

Bae, Y. J., 278
Bahlai, C. A., 50
Bai, Y. Y., 243
Baker, T. C., 169
Bakker, J. D., 1159
Balciunas, J., 114, 841
Barber, N. A., 492
Barraclough, E. I., 1359
Barrett, J. P., 633
Barro, P., 1652
Basoalto, E., 1399
Bauce, E., 498
Bauer, L. S., 1513
Bauman, T. A., 633
Bean, D. W., 1666
Beers, E. H., 286
Belovsky, G. E., 312
Bencivenga, N., 1170
Bennett, K. C., 1631
Bento, F.d.M.M., 1315
Bento, J.M.S., 678
Berenbaum, M.R., 1291, 1922
Berkebile, D. R., 1101
Bethke, J. A., 159
Beuzelin, J. M., 528
Bi, M.-J., 883
Biddinger, D. J., 2045
Björklund, N., 898
Blair, D., 1122
Blake, A. J., 1195
Blankenship, E. E., 202
Bledsoe, L., 484
Boetel, M. A., 1545
Bosch, J., 149
Botha, A.-M., 1206
Branson, D. H., 346, 1441
Breidenbaugh, M. S., 591
Brewer, M. J., 31
Brightwell, R.j., 1659
Broadbent, A. B., 1570
Broce, A. B., 1101
Brock, C., 1961
Brousseau, P.-M., 1151
Brown, G. C., 1309, 1788
Brown, J. K., 1028
Brown, M. D., 545
Brown, M. W., 134
Buczkowski, G., 1936
Bueno, V.H.P., 1500
Burgess, E.P.J., 1359
Butler, R. C., 303
Byun, B.-K., 337

C

Cafaro, M. J., 105
Calatayud, P.-A., 57
Campbell, J. B., 1101
Campbell, J. F., 919
Campos, M., 79
Candia, O., 1276
Cao, W.-J., 232
Carneiro, M.A.A., 849, 1134
Carney, V. A., 738, 1666
Carpaneto, G. M., 1756
Carpenter, J. E., 1884
Carreira, V. P., 865
Carver, S., 798
Cavanagh, A. F., 1953
Chaŕez-Moreno, C. K., 1178
Chabi-Olaye, A., 468
Chacón, E., 1848
Chapman, R. B., 303
Chen, H.-S., 1021
Chen, M., 715, 1369
Chen, P., 1378
Chen, W., 1705
Chen, Wenlong., 1545
Chen, X. X., 1369
Chen, X.-X., 2039
Chen, Y., 715, 1369
Cheng, D.-F., 1737
Cheng, J. A., 243
Cheng, R.-X., 389
Cherry, R. H., 1012, 2025
Chinnasamy, D., 1980
Chiozza, M. V., 856
Choate, B., 1428, 1447
Chu, D., 1028
Chun, J. H., 337
Clayton, M. K., 874
Cloutier, C., 1151
Cockfield, S. D., 286
Cokendolpher, J. C., 1428
Collins, J. A., 1428
Colunga-Garcia, M., 31
Cook, S. P., 821
Coop, L. B., 2006
Cooper, W. R., 223, 787
Corcuera, P., 320
Corio, C., 865
Corn, J. G., 164
Cornelissen, T., 849
Cory, J. S., 459
Costa, C.M.Q., 359
Costa, M.l.Z., 1315
Costa-Lima, T. C., 195
Cotes, B., 79
Cottrell, T. E., 610
Coyle, D. R., 1242, 1539
Currie, C. R., 105, 406
Curtis, P. S., 1810

D

Daane, K. M., 1232, 1264
Dale, P., 88
Dalin, P., 1666
Daniels, J. C., 1961
Danne, A., 970
Davidson, L. N., 576
Davis, A.L.V., 811
Davis, G. A., 440
Debano, S. J., 1
Defoliart, L. S., 1643
Delatte, J., 1715
Delpuech, J. M., 535
Delucia, E. H., 1291
Deschodt, C. M., 811
De M.M. Bento, F., 1315
De La Mora, A., 1473
De Vries, R. S. M., 459
De Szalay, F. A., 591
De Lyra, M. C. C. P., 1178
Dickey, A. M., 1037
Dieng, H., 1409
Dillwith, J. W., 223
Dively, G. P., 707
Donaldson, J., 1561
Dorn, S., 561

Dosdall, L. M., 1092, 1195
Drayton, G. M., 303
Dreves, A. J., 2006
Drummond, F. A., 1428, 1447
Duan, J. J., 1513
Dudley, T. L., 1666
Dunkel, F. V., 979
Dupas, S., 57
Dupont, C., 535
Dutra, V. S., 1457

E

Eberts, D., 1666
Eckhardt, L. G., 415
Edde, P. A., 930
Eliopoulos, P. A., 1352
Elliott, N. C., 763
Endo, N., 1973
Epsky, N. D., 1506
Erasmus, A., 702
Esteves, D., 849
Evans, E. W., 576
Evens, T. J., 670

F

Fang, Q., 715
Farhadi, R., 1586
Farias, A. I., 359
Fathipour, Y., 177
Favi, F. D., 351
Favila, M. E., 328
Feagin, R. A., 127
Feken, M. A., 1961
Feng, H., 1378, 1698
Fergen, J. K., 1816
Fernandes, G. W., 849, 1134
Fielding, D. J., 1643
Fierke, M. K., 520
Flaherty, L., 1794
Fleischer, S. J., 2045
Floreani, C., 1652
Follum, R. A., 1858
Fombong, A. T., 1731
Fork, S. K., 753
Forschler, B. T., 98
Forsythe, Jr., H. Y., 1428
Frank, J. H., 939
Fraser, I., 253
Frewin, A. J., 1570
Fuchs, M., 266
Fuentes-Contreras, E., 1399

G

Gao, M.-Q., 2039
Gao, P., 459
Gao, Xiwu, 600
Gao, Xue, 1878
Gao, Y., 1698
Gara, R. I., 1159
Garcia-Marí, F., 827
Gardner, K. T., 1666
Gava, C., 570
Ge, F., 2033
Georgievska, L., 459
Gerard, P. D., 1420
Geremias, L. D., 195
Ghimire, M. N., 449
Gilbert, R. A., 1012, 2025
Giles, K. L., 763
Gillette, N. E., 406
Giordano, R., 1170
Glas, J. J., 1922
Godfrey, L. D., 685, 1255, 1617
Gomez-Torres, M. L., 678
Gontijo, L. M., 286
Gould, J., 1513
Grant, J. F., 1858, 1866
Gratton, C., 31, 1561
Greene, J. K., 956, 1420
Groden, E., 1447
Groves, R. L., 1264
Gu, L.-Q., 1249
Guédot, C., 1302
Guimarães, J. A., 1457
Guo, J.-Y., 1021
Guo, Y., 600

H

Hébert, C., 1151
Haavik, L. J., 520
Haddad, M. L., 678
Hagen, E. N., 1159
Halffter, G., 1165
Hall, D. G., 670
Hallett, R. H., 50, 1570
Hamm, P. B., 1
Hammack, L., 727
Hamud, S. M., 642
Han, E., 498
Han, Y., 1929
Hanna, R., 695
Hao, L., 1929
Harris, M. K., 1593
Hasson, E., 865
Hayes, J. L., 1685
Haynes, K. F., 1309
Hazzard, R. V., 1953
Heard, S. B., 1794
Hebertson, E., 1666
Heckel, D. G., 552
Heimpel, G. E., 31
Hernández-Delgado, S., 1043
Hoepting, C. A., 266
Hoffmann, A. A., 970
Hofs, J.-L., 1523
Hogsette, J. A., 1101
Holdcraft, R., 1980
Hong, S. C., 1561
Hong, X.-Y., 1801
Hoosier, B. M., 633
Hoover, K., 169
Horton, D. R., 1302
Hou, M., 1344, 1929
Hou, S.-P., 2039
Hou, Y., 1676
Hribar, L. J., 1961
Hsu, C. L., 266
Hu, C., 715, 1369
Hu, G., 1705
Hu, W., 88
Huang, C.-Y., 1705
Huang, F., 1737
Huang, F. N., 243
Huang, S.-C., 999
Huerta, C., 1165
Hunt, T. E., 202
Hurst, T. P., 545
Husseneder, C., 378, 1715
Hwang, S.-Y., 1990

I

Ieradi, M., 1756

J

Jackson, R. E., 1085, 1111
Jadhav, R., 892
Jandricic, S. E., 1631
Jaronski, S. T., 979
Jennings, D. T., 1428
Jensen, A., 1
Jensen, P. D., 707
Jezorek, H. A., 1884
Jiang, X., 1579
Jiang, X.-F., 232
Jiménez-Salinas, E., 320
Johnson, Douglas W., 1309
Johnson, D. W., 1788
Johnson, M. T., 1848
Johnson, M. W., 1264
Jones, E. N., 738, 1666
Jones, G. V., 2006
Jones, S. C., 1810
Jordan, M. S., 1539
Juliano, S. A., 1586

K

Kairu, E. W., 57
Karlovsky, P., 1532
Kaufman, L. V., 430
Kaufman, P. E., 1608
Kay, B. H., 545
Kazmer, D. J., 1666
Kean, A. M., 1359
Keddie, A., 1092
Keddie, B. A., 1195
Keena, M. A., 169, 1323
Kells, S. A., 661
Khanjani, M., 177
Khrimian, A., 610
Kim, H., 278
Kim, J. G., 278
Kim, S.-S., 337
Klassen, W., 1389
Klein, A.-M., 1744
Kline, D. L., 1608
Klingenberg, M. D., 898

Knight, A. L., 583, 1283, 1399
Knolhoff, L. M., 1922
Kok, L. T., 440
Kong, H., 1579
Kontodimas, D. C., 1352
Kowles, K. A., 892
Kraemer, M. E., 351
Krogh, P. H., 1170
Krugner, R., 1264
Kryger, U., 811
Kubota, K., 1336
Kuenen, L.P.S., 1059
Kumar, V., 1389
Kurtz, B., 1532
Kwon, T.-S., 337

L

López, M. A., 295
López-Ortega, G., 320
López-Portillo, J., 1165
Labadie, P.e., 1659
Lambdin, P. L., 1858, 1866
Lamp, W. O., 707
Lampinen, B. D., 1232
Landolt, P. J., 1, 1302
Lapitan, N.L.V., 1206
Lapointe, S. L., 670
Laws, A. N., 312
Leblanc, L., 989, 1626
Leclair, G., 1997
Lee, C.-Y., 835
Lee, J. C., 642, 653
Legaspi, Jr., B. C., 368
Legaspi, J. C., 368
Lemay, M. A., 1997
Leopold, R. A., 1545
Leppla, N. C., 939
Leslie, T. W., 2045
Lewis, J. L., 98
Li, B.-P., 389
Li, G., 1378, 1698
Li, H.-F., 1601
Li, L., 999
Li, Q.-L., 883
Li, Y., 1051
Liao, X., 1929
Lill, J. T., 1893
Lim, J.-H., 337
Lin, Z.-X., 1705
Lindgren, B. S., 476
Lins, Jr., J. C., 1500
Liu, B., 1378
Liu, C., 1051
Liu, H., 1737
Liu, T.-X., 883, 912
Liu, X., 1878
Liu, Y., 1344
Liu, Z., 552
Liu, Z. C., 1369
Lizama, M., 1276
Lobo, J. M., 42
Lopes, F.S.C., 570
Luckhart, S., 396
Lundgren, J. G., 1816
Luo, L., 1579
Luo, L.-Z., 232
Lv, J., 528
Lyra, M.C.C.P.D., 1178

M

Ma, Z.-L., 999
MacIntosh, G. C., 856
Mafra-Neto, A., 1980
Magee, D., 1997
Mahroof, R. M., 930
Mailafiya, D. M., 57
Maistrello, L., 295
Makino, S., 424
Malavasi, A., 570
Mallis, R. E., 1466
Malone, L. A., 1359
Manen, F.T.v., 1866
Manfrin, M., 865
Maniania, N. K., 468
Mannion, C. M., 1506
Manosalva, L., 1276
Marcum, D. B., 685, 1617
Marques, R. N., 1315
Marshall, D. B., 625
Martínez-Romero, E., 1178
Matusick, G., 415
Mattson, Jr., W. J., 1242
Mauffette, Y., 498
Mausel, D. L., 440
Mayek-Pérez, N., 1043
Mazziotta, A., 1756
Mccullough, D. G., 253
Mcinnis, D., 570
McSorley, R., 939
Medina, R. F., 1037
Mehelis, C., 114, 841
Meiler, S. U., 979
Men, X., 2033
Meng, L., 389
Mengersen, K., 88
Meuti, M. E., 1810
Michels, Jr., C. J., 738, 1666
Migiro, L. N., 468
Milbrath, L. R., 68
Miller-Pierce, M. R., 513
Mills, N. J., 1232
Miranda, M., 1399
Mironidis, G. K., 1068
Moharramipour, S., 177
Montá, L. D., 1652
Moon, R. D., 1101
Moore, P. M., 1323
Mora, A.d.l., 1473
Morelli, R., 570
Morgan, D.J.W., 1264
Morse, J. G., 15
Moura, R. C., 359
Mueller, E. E., 31
Murphy, S. M., 1893
Mutis, A., 1276
Muto, S., 1973
Muzika, R. M., 1482

N

Nam, J. M., 278
Nathan, B. P., 892
Nault, B. A., 266
Nava, D. E., 678
Nechols, J. R., 919
Negrón, J. F., 642
Nehme, M. E., 169
Neoh, K.-B., 835
Neser, S., 841
Newman, J., 50
Nicolai, N., 127
Niedz, R. P., 670
Nielsen, M.-C., 303
Nink, R. A., 1457
Noma, T., 31
Nor, S.A.M., 210
Notter-Hausmann, C., 561
Nuessly, G. S., 1012, 2025

O

Obenauer, P. J., 1608
Ocete, R., 295
Okabe, K., 424
Olfert, O., 1092
Oliveira, P. S., 1903
Olson, B. E., 775
O'meara, S. A., 1666
O'Neal, M. E., 31, 856
O'Neil, R. J., 484
O'Neill, B. F., 1291
O'Neill, K. M., 775
Onzo, A., 695
Opit, G. P., 1004
Orwig, D. A., 513
Ouellette, G. D., 1447

P

Pérez-Hernández, M. A., 320
Palma, R., 1276
Palmer, C. M., 1775
Papadopoulos, N. T., 570
Paranhos, B. J., 570
Pardo, F., 1276
Parra, J.R.P., 195, 678, 1315
Parra, L., 1276
Pascual, F., 79
Patt, J. M., 618
Pavan, F., 1652
Payton, M. E., 1004
Pekas, A., 827
Penfold, C. M., 970
Peng, Y. F., 715, 1369
Peng, Z.-Q., 999
Perera, O. P., 1111
Perich, F., 1276
Pfeiffer, D. G., 396
Philip, B. A., 1359

Philipsen, H., 303
Phillips, T. W., 449, 930
Philpott, S. M., 1473
Phoofolo, M. W., 763
Pierce, S. J., 253
Pierre, J., 1523
Pikul, Jr., J. L., 727
Pitts-Singer, T. L., 149
Poland, T. M., 253
Polk, D., 1980
Pollock, D. A., 738
Portman, S. L., 939
Potter, D. A., 1829, 1838
Powell, B. E., 1492
Powell, J. S., 874
Preisser, E., 513
Pu, D.-Q., 2039
Puckett, R. T., 1593
Puckette, J. A., 930
Puterka, G. J., 223

Q

Qin, W.-Q., 999
Qiu, F., 1378
Quiroz, A., 1276

R

Raffa, K. F., 406, 874, 1242, 1539
Ramírez-Puebla, S. T., 1178
Ranney, J. W., 1858, 1866
Rao, S., 2017
Rawi, C.S.M., 210
Reagan, T. E., 528
Reay-Jones, F.P.F., 944, 956, 1420, 1943
Redak, R. A., 159
Redmond, C. T., 1838
Reed, A., 1858
Reed, S. E., 1482
Reeves, R. B., 956, 1420
Reichert, E., 1848
Reisig, D. D., 685, 1255, 1617
Reyna, S. M., 1037
Rhainds, M., 484, 1794
Richards, S. L., 190
Rieske, L. K., 787, 1466
Riley, D. G., 505
Rinehart, J. E., 1685
Robert, J. A., 476
Robertson, B., 930
Rochefort, S., 498
Rodrigues, M. D., 678
Rodriguez-Saona, C., 1980
Rohr, J. R., 2045
Rolston, M. G., 775
Romero, S. A., 919
Rondon, S. I., 1, 1037
Rosas-García, N. M., 1043
Rosenblueth, M., 1178
Rowles, A. D., 1141
Ru, B.P.L., 57
Ruano, F., 79
Rubinoff, D., 989, 1626
Ruiz, E. A., 1685
Ryall, K. L., 121
Ryan, P. A., 545

S

Sétamou, M., 618
Sabelis, M. W., 695
Sadof, C. S., 484
Salcedo, C., 907
Salmah, M.R.C., 1409
Salom, S. M., 440
Salomão, A. T., 1724
Sampaio, M. V., 1500
Sanderson, J. P., 1631
Sandhu, H. S., 1012, 2025
Santos, M. S., 1457
Sarmiento-Benavides, S. L., 1043
Sasakawa, K., 1554
Sasaki, R., 1973
Sasu, M. A., 140
Saunders, M. C., 1751
Savopoulou-Soultani, M., 1068
Schaafsma, A. W., 50, 1570
Scheirs, J., 552
Scherr, M. A., 2017
Schnitzler, F.-R., 1359
Scholl, P. J., 1101
Scholtz, C. H., 811
Schwartzberg, E. G., 1309, 1788
Seal, D. R., 1389
Sedlak, C. W., 979
Seibert, C. E., 775
Seidl-Adams, I., 140
Seybold, S. J., 642, 661
Sharley, D. J., 970
Shelton, A. M., 266
Shepherd, W. P., 633
Shi, M., 2039
Shi, P., 2033
Shin, J. H., 337
Shirley, B. M., 821
Short, B. D., 134
Sidney, L. A., 1500
Siegel, J. P., 1059
Siegert, N. W., 253
Sikkema, S., 50
Silk, P. J., 1794, 1997
Silva, A. P., 1315
Silva, D. B., 1500
Silva, D. P., 1903
Silva, F.A.B., 359
Silva, J. G., 1457
Silva, N.M.O., 1457
Silverman, J., 1141, 1492, 1659
Simms, D. M., 1715
Skinkis, P. A., 2006
Smartt, C. T., 190
Smeins, F. E., 127
Snodgrass, G. L., 1085, 1111
Son, Y., 1264
Sorensen, M. A., 159
Soto, E. M., 865
Soto, I. M., 865
Soto-Adames, F., 1170
Sousa-Souto, L., 1744
Spafford, H., 798
Spencer, J. L., 1922
Stamopoulos, D. C., 1068
Stathas, G. J., 1352
Stavrinides, M. C., 1232
Stephen, F. M., 520
Stephenson, A. G., 140
Stiling, P. D., 1884
Stilwell, A. R., 202
Storey, A., 798
Story, J. M., 164
Strümpher, W. P., 811
Su, N.-Y., 1601
Sullivan, B. T., 633
Sun, X., 459
Swan, C. M., 707
Swanevelder, Z. H., 1206
Sweeney, J. D., 1794, 1997
Switzer, P. V., 892
Sword, G. A., 1441
Szalay, F.A.D., 591

T

Taerum, S. J., 105
Tan, C.-W., 1990
Tansey, J. A., 1092
Taylor, D. B., 1101
Teal, P.E.A., 1731
Tecante, A., 1178
Telhado, C., 849
Tena, A., 827
Teodoro, A. V., 1744
Teulon, D.A.J., 303
Thompson, D. C., 1666
Thomson, L. J., 970
Throne, J. E., 1004
Tian, J. C., 715, 1369
Tigreros, N., 892
Tillman, P. G., 610, 1184, 1765
Timer, J., 1751
Tobin, P. C., 1751
Toews, M. D., 956, 1420
Toki, W., 1336
Tong, S., 88
Torto, B., 1731
Trager, M. D., 1961
Trimble, R. M., 625
Trotta-Moreu, N., 42
Trudeau, M., 498
Tscharntke, T., 1744
Tu, Z., 396

U

Ulyshen, M. D., 1513

V

Valverde, P. L., 320
Vandenberg, J., 468, 702

VanManen, F. T., 1866
Vander Werf, W., 459
Van Driesche, R. V., 1513
Vanek, S. J., 1829
Vanweelden, M., 1936
Vargas, R. I., 989, 1626
Vasconcellos-Neto, J., 1724
Venette, R. C., 661
Veo, K. D., 979
Vidal, D. B., 1457
Vidal, S., 1532
Villegas-Mendoza, J. M., 1043
Vite, F., 320
Vitek, C. J., 190
Vlak, J. M., 459

W

Waite, J., 1170
Walder, J.M.M., 1315
Wall, K., 140
Wallander, R., 775
Walter, A. J., 661
Walter, C., 1359
Walton, V. M., 2006
Wan, F.-H., 1021, 1028
Wang, C.-X., 883
Wang, C.-Z., 1249
Wang, J.-J., 1737
Ward, J. S., 1911
Webb, S. E., 1012, 2025
Weber, E., 1170
Weinstein, P., 798
Welsman, J. A., 1570
Weng, Z., 1676
West, M. S., 727
Westhuizen, L.V.D., 841
Westwood, A. R., 1122
Wheeler, T. A., 1848
White, L. J., 164
Wiggins, G. J., 1858, 1866
Wilkerson, J. B., 1858, 1866
Williams, D. W., 253
Williams, S. C., 1911
Wilson, L. T., 528
Winsor, J. A., 140
Witcosky, J. J., 642
With, K. A., 919
Woodman, D., 1428
Wooster, D. E., 2017
Wraight, S. P., 1631
Wright, M. G., 430
Wright, R. J., 202
Wu, K., 600, 1051, 1344, 1698
Wu, S., 1378
Wyckhuys, K.A.G., 31, 600, 1698

X

Xie, M.-C., 1705
Xin, D.-Y., 1705
Xu, Z., 169
Xue, M., 883
Xue, Y., 1570

Y

Yadav, J., 1990
Yan, R. H., 243
Yan, W., 999
Yan, X.-H., 1737
Yang, R.-L., 1601
Yaninek, S., 484
Ye, G. Y., 243, 715, 1369, 2039
Yoo, H.J.S., 484
Yoon, J., 278
Yu, H., 600

Z

Zahiri, B., 177
Zalucki, M. P., 177
Zambino, P. J., 821
Zandigiacomo, P., 1652
Zang, L.-S., 912
Zangerl, A. R., 1291
Zanzot, J. W., 415
Zhai, B.-P., 1705
Zhang, A., 169
Zhang, H., 1801
Zhang, J.-H., 1249
Zhang, K.-J., 1801
Zhang, L., 232, 1579
Zhang, X., 396
Zhang, X.-X., 1705
Zhang, Y., 1051
Zhang, Y. J., 1028
Zhang, Y., 600
Zhao, Z.-M., 1737
Zhong, H., 1961
Zhou, Z.-S., 1021
Zuñiga, G., 1685

CUMULATIVE KEY WORD INDEX
Environmental Entomology
Volume 39, 2010

A

α-bisabolene epoxide,
  brown stink bug, 610
  green stink bug, 610
  methyl (*E,E,Z*)-2,4,6-decatrienoate, 610
  southern green stink bug, 610
abiotic factors,
  fecundity, 195
  vegetable leafminer, 195
abiotic interactions,
  environmental limitation, 1659
  invasive species, 1659
  low temperature, 1659
  range limitation, 1659
above- and below-ground interactions,
  indirect effects, 1561
  induced plant responses, 1561
  phytoparasite, 1561
  root herbivory, 1561
absolute sampling,
  integrated pest management, 1617
  relative sampling, 1617
  wing dimorphism, 1617
  wing diphenism, 1617
Acadia National Park,
  Formicidae, 1447
  Myrmica rubra, 1447
  species, 1447
  survey, 1447
Acalymma vittatum, 140, 1953
  bacterial wilt disease, 140
  butternut squash, 1953
  Cucurbita maxima, 1953
  Cucurbita pepo ssp, 140
  integrated pest management (IPM), 1953
  pollination, 1953
  real-time polymerase chain reaction, 140
  texana, 140
  Erwinia tracheiphila, 140
acarinarium,
  Eumeninae, 424
  guarding mutualism, 424
  host control, 424
  parasite, 424
Acer saccharum,
  exotic species, 1539
  folivore, 1539
  plant-insect interactions, 1539
  Rubus, 1539
Acrididae,
  global warming, 1643
  insect life cycles, 1643
  Orthoptera, 1643
  voltinism, 1643
Adelges tsugae, 440, 513
  establishment, 440
  Fiorinia externa, 513
  herbivory, 513
  Laricobius nigrinus, 440
  release strategy, 440
  Tsuga canadensis, 440, 513
Adelphocoris suturalis Jakovlev,
  nontarget pest, 1378
  oviposition behavior, 1378
  population density, 1378
  transgenic cotton, 1378
adult emergence,
  Mangifera indica, 1409
  population dynamics, 1409
  thrips, 1409
adulticides,
  biting midges, 591
  Diptera, 591
  Hymenoptera, 591
Aegorhinus superciliosus,
  behavioral bioassay, 1276
  electroantennogram, 1276
  R-limonene, 1276
aerial ULV application,
  Miami blue butterfly, 1961
  mosquito control, 1961
  naled, 1961
  nontarget impact, 1961
Aethina tumida,
  honey bee, 1731
  Kenya, 1731
  Kodamaea ohmeri, 1731
  Pheidole megacephala, 1731
aggregation, 892, 1420, 1794
  Allee effect, 1794
  competitive displacement, 1794
  dispersion, 1420
  edge effect, 1420
  fecundity, 892
  Japanese beetle, 892
  lipid, 892
  mate choice, 892
  SADIE analysis, 1794
  spatial-temporal variation, 1420
  spatial association, 1794
aggregation pheromone, 642, 874
  kairomones, 642
  Scolytidae, 642
  Scolytus multistriatus, 642
  Scolytus schevyrewi, 642
  Ips pini, 874
  ipsdienol, 874
  monoterpenes, 874
  variance components, 874
agonistic behavior,
  invasive species, 1601
  subterranean termite, 1601
  tunnel blockage, 1601
Agrilus planipennis,
  biological invasion, 253
  dendroentomology, 253
  emerald ash borer, 253
  eradication, 253
agroforest,
  biodiversity, 1473
  Eucharitidae, 1473
  poneromorph, 1473
  trophic interaction, 1473
Agrotis segetum,
  Bt maize, 702
  Cry1Ab, 702
  cutworm, 702
air movement,
  habitat quality, 919
  movement behavior, 919
  shelter, 919
  Tribolium castaneum, 919
alfalfa insects,
  alfalfa management, 202
  degree-day, 202
  Hypera postica, 202
alfalfa leafcutting bee,
  bee density, 149
  Medicago sativa, 149
  reproductive success, 149
alfalfa management,
  alfalfa insects, 202
  degree-day, 202
  Hypera postica, 202
alfalfa weevil,
  development time, 177
  immature survival, 177
  modeling, 177
  rate isomorphy, 177
Allee effect,
  aggregation, 1794
  competitive displacement, 1794
  SADIE analysis, 1794
  spatial association, 1794

Allium cepa,
epidemiology, 266
management, 266
Thrips tabaci, 266
Tospovirus, 266
almonds,
navel orangeworm, 1059
overwintering emergence, 1059
pistachios, 1059
sex ratio, 1059
Amara,
copulatory behavior, 1554
feeding ecology, 1554
Harpalus (*Pseduoophonus*), 1554
predispersal seed predation, 1554
Amazon,
biodiversity, 1134
galling insects, 1134
mining, 1134
restoration ecology, 1134
Amblydromalus manihoti,
cassava green mite, 695
Euseius fustis, 695
Phytoseiidae, 695
Typhlodromalus aripo, 695
Ambrosia artemisiifolia,
development, 1021
fecundity, 1021
Ophraella communa, 1021
temperature, 1021
American chestnut,
chestnut weevil, 787
Chinese chestnut, 787
Ormyrus labotus, 787
Torymus sinensis, 787
ammonium,
attractant, 989
Drosophilidae, 989
nontarget, 989
putrescine, 989
ampicillin,
gut flora accommodation, 1170
oxytetracycline, 1170
tylosin, 1170
Wolbachia, 1170
amplified fragment length polymorphism, 232, 1037
gene flow, 232
geographic population, 232
Loxostege sticticalis, 232
migration, 232
phylogeography, 1037
potato, 1037
potato tubermoth, 1037
amplified fragment length polymorphism markers,
genetic structure, 1043
insect pest, 1043
pink hibiscus mealybug, 1043
polymorphism, 1043
Anagrus nilaparvatae,
developmental time, 2039
fecundity, 2039
successive generation, 2039
transgenic Bt rice, 2039
Anastrepha,
hosts, 1457
parasitoids, 1457
Tephritidae, 1457
ANCOVA,
Chow test, 2033
hypothesis of isomorphy rate, 2033
lower developmental threshold, 2033
androcyclic mode,
anholocyclic mode, 1878
Aphis gossypii, 1878
cotton, 1878
intermediate mode, 1878
anholocyclic mode,
androcyclic mode, 1878
Aphis gossypii, 1878
cotton, 1878
intermediate mode, 1878
Anomala (=*Exomala*) *orientalis*,
(Z)-7-tetradecen-2-one, 1980
blueberries, 1980
hand-applied plastic dispensers, 1980
Anoplophora glabripennis,
male-produced pheromone, 169
monitoring traps, 169
plant volatiles, 169
ant activity,
biological control, 827
Lasius grandis, 827
Pheidole pallidula, 827
Plagiolepis schmitzii, 827
ant nest-seedling spatial relationship,
grassland spatial structure, 127
Pogonomyrmex barbatus, 127
Ripley's K function, 127
seedling establishment, 127
ant-tending,
Aphis glycines, 1309
Lasius neoniger, 1309
Scymnus louisianae, 1309
antenna,
Choristoneura rosaceana, 625
Grapholita molesta, 625
pheromone, 625
sensory adaptation, 625
antibiosis,
Aphis glycines resistance, 856
free amino acids, 856
ant–aphid interaction,
Argentine ant, 1492
cotton aphid, 1492
green peach aphid, 1492
odorous house ant, 1492
ants, 1788, 1903
aphid-tending, 1788
Aphis glycines, 1788
Caryocar brasiliense, 1903
cerrado savanna, 1903
Lasius neoniger, 1788
myrmecophily, 1903
stink bugs, 1903
aphid diets,
aphid saliva collection, 223
Russian wheat aphid, 223
saliva proteins, 223
Triticum aestivum, 223
aphid saliva collection,
aphid diets, 223
Russian wheat aphid, 223
saliva proteins, 223
Triticum aestivum, 223
aphid-tending,
ants, 1788
Aphis glycines, 1788
Lasius neoniger, 1788
aphidiidae,
Buchnera aphidicola, 389
development, 389
endosymbiosis, 389
life history, 389
Aphidius ervi,
biological control, 1500
larval competition, 1500
Macrosiphum euphorbiae, 1500
Praon volucre, 1500
aphids,
biological control, 763
Coccinellidae, 763
relay-intercropping, 763
sorghum, 763
Aphis fabae,
functional response, 1586
handling time, 1586
Hippodamia variegata, 1586
searching efficiency, 1586
Aphis glycines, 31, 50, 1309, 1788
ants, 1788
ant-tending, 1309
aphid-tending, 1788
biological control, 31
hedgerow, 50
landscape ecology, 31
Lasius neoniger, 1309, 1788
model selection, 50
plant chemistry, 31
potassium, 31
Rhamnus cathartica, 50
Scymnus louisianae, 1309
Aphis glycines resistance,
antibiosis, 856
free amino acids, 856
Aphis gossypii,
androcyclic mode, 1878
anholocyclic mode, 1878

cotton, 1878
intermediate mode, 1878
Apis mellifera,
Astylus atromaculatus, 1523
Bt cotton, 1523
gene flow, 1523
insect population, 1523
apple, 134, 583, 1283
codlemone, 1283
Cydia pomonella, 583, 1283
lures, 583
mating disruption, 1283
monitoring, 583
stink bug, 134
traps, 1283
UV radiation, 134
wound response, 134
arboreal foraging,
Argentine ants, 1141
biological invasion, 1141
Linepithema humile, 1141
Pinus taeda, 1141
areawide,
Cydia pomonella, 1399
geostatistics, 1399
immunomarking, 1399
mark-capture, 1399
Argentine ant, 1141, 1492
ant–aphid interaction, 1492
arboreal foraging, 1141
biological invasion, 1141
cotton aphid, 1492
green peach aphid, 1492
Linepithema humile, 1141
odorous house ant, 1492
Pinus taeda, 1141
arthropod communities,
Juan Fernández Archipelago, 1159
plant invasion, 1159
trophic levels, 1159
arthropod damage,
hyperspectral, 1255
multispectral, 1255
artificial injury,
compensation, 528
rice, 528
sugarcane borer, 528
Asian longhorned beetle,
degree-days, 1323
development, 1323
pupation, 1323
survival, 1323
temperature, 1323
Asian tiger mosquito,
aspirator, 1608
BG-Sentinel, 1608
landing counts, 1608
surveillance, 1608
Asiatic rice borer,
host orientation, 1929
host range, 1929
Lepidoptera, 1929
olfaction, 1929
aspirator,
Asian tiger mosquito, 1608
BG-Sentinel, 1608
landing counts, 1608
surveillance, 1608
assemblage structure,
dung, 811
Kalahari, 811
landscape, 811
Tswalu, 811
Astylus atromaculatus,
Apis mellifera, 1523
Bt cotton, 1523
gene flow, 1523
insect population, 1523
Atlantic rainforest,
clear-cut area, 359
community structure, 359
dung beetle ecology, 359
atmospheric nitrogen,
isotope discrimination factor, 1810
stable isotope ratios, 1810
termites, 1810
attractant, 633, 989
ammonium, 989
bark beetle pheromone, 633
Drosophilidae, 989
endo-brevicomin, 633
exo-brevicomin, 633
gas chromatography electroantennographic detection, 633
nontarget, 989
putrescine, 989
Aulacorthum solani,
developmental rate, 1631
foxglove aphid, 1631
greenhouse floriculture pests, 1631
Lactin model, 1631
autoinoculation device,
Beauveria bassiana, 468
leafminer, 468
Liriomyza huidobrensis, 468
Metarhizium anisopliae, 468

B

B biotype-specific primers,
Bemisia tabaci, 1028
displacement, 1028
invasion biology, 1028
Bacillus thuringiensis, 707, 1369
Bt rice, 1369
genetically modified crops, 707
nontarget effect, 1369
nontarget effects, 707
stream ecology, 707
tritrophic interactions, 1369
Ummeliata insecticeps, 1369
bacteria,
cellulase, 378
gut, 378
protozoa, 378
16S rRNA gene, 378
bacterial wilt disease,
Acalymma vittatum, 140
Cucurbita pepo ssp, 140
real-time polymerase chain reaction, 140
texana, 140
Erwinia tracheiphila, 140
bacteriocytes,
Coccoidea, 1178
endosymbiont, 1178
scale insects, 1178
systematics, 1178
Bactrocera,
BioLure, 1626
Ceratitis, 1626
nontarget, 1626
torula yeast, 1626
baited pitfall traps,
beetles, 1151
coprophilous, 1151
forest types, 1151
necrophagous, 1151
bark beetle, 406, 415, 661
boring incitant, 661
community analysis, 406
host-range prediction, 661
invasive species, 661
Leptographium spp, 415
microbial diversity, 406
Ophiostomatoid fungi, 415
Pinaceae, 661
Pinus palustris, 415
regeneration weevils, 415
symbiosis, 406
bark beetle pheromone,
attractant, 633
endo-brevicomin, 633
exo-brevicomin, 633
gas chromatography-electroantennographic detection, 633
Beauveria bassiana,
autoinoculation device, 468
leafminer, 468
Liriomyza huidobrensis, 468
Metarhizium anisopliae, 468
bee density,
alfalfa leafcutting bee, 149
Medicago sativa, 149
reproductive success, 149
beetles,
baited pitfall traps, 1151
coprophilous, 1151
forest types, 1151
necrophagous, 1151

behavior,
locomotor activity, 1922
oviposition, 1922
rotation resistance, 1922
soybean herbivory, 1922
behavioral bioassay,
Aegorhinus superciliosus, 1276
electroantennogram, 1276
R-limonene, 1276
behavioral experiments,
gas chromatography–electroantennographic detection, 600
Helicoverpa armigera, 600
Microplitics mediator, 600
plant volatiles, 600
behavioral trade-offs,
Helicoverpa armigera, 552
larval performance, 552
oviposition preference, 552
realized fecundity, 552
belowground interactions,
Diabrotica virgifera virgifera, 1532
Fusarium verticillioides, 1532
larval development, 1532
root colonization, 1532
Bemisia tabaci, 912, 1028
B biotype-specific primers, 1028
displacement, 1028
Encarsia, 912
Eretmocerus, 912
invasion biology, 1028
sweetpotato whitefly, 912
Berberis thunbergii,
Borrelia burgdorferi, 1911
Ixodes scapularis, 1911
microclimate, 1911
vapor pressure deficit, 1911
BG-Sentinel,
Asian tiger mosquito, 1608
aspirator, 1608
landing counts, 1608
surveillance, 1608
bioclimatic models,
Cactoblastis cactorum, 368
cactus moth, 368
climate envelope, 368
model validation, 368
biodiversity, 1134, 1473, 1816
agroforest, 1473
Amazon, 1134
biological control, 1816
conservation, 1816
Elymus trachycaulus, 1816
Eucharitidae, 1473
galling insects, 1134
generalist predators, 1816
IPM, 1816
mining, 1134
poneromorph, 1473
restoration ecology, 1134
trophic interaction, 1473
western corn rootworm, 1816
biological control, 31, 430, 449, 763, 827, 1352, 1500, 1506, 1570, 1666, 1816, 1848, 1858
ant activity, 827
Aphidius ervi, 1500
aphids, 763
Aphis glycines, 31
biodiversity, 1816
biological invasions, 1666
Carduus nutans, 1858
Chilocorus bipustulatus, 1352
Cirsium, 1858
Coccinellidae, 763
conservation, 1816
Costa Rica, 1848
critical daylength response, 1666
development, 1352
Elymus trachycaulus, 1816
functional response, 1570
generalist predators, 1816
Hawaiian archipelago, 430
host suitability, 449
host-specificity, 1848
hosts, 1506
invasive species, 1848
IPM, 1816
landscape ecology, 31
larval competition, 1500
Lasius grandis, 827
life history, 1848
Macrosiphum euphorbiae, 1500
nontarget, 430, 1858
parasitism, 1506
parasitism rates, 430
parasitoid, 449, 1506
Pheidole pallidula, 827
Plagiolepis schmitzii, 827
plant chemistry, 31
potassium, 31
Praon volucre, 1500
relay-intercropping, 763
Rhinocyllus conicus, 1858
sorghum, 763
stored-product insects, 449
temperature, 1352
temperature-dependent development, 1570
thermal constant, 1352
voltinism, 1666
wasp, 449
western corn rootworm, 1816
biological invasion, 253, 1141, 1666, 1884
Agrilus planipennis, 253
arboreal foraging, 1141
Argentine ants, 1141
biological control, 1666
cactus moth, 1884
critical daylength response, 1666
dendroentomology, 253
emerald ash borer, 253
eradication, 253
larval performance, 1884
Linepithema humile, 1141
Opuntia, 1884
oviposition preference, 1884
Pinus taeda, 1141
voltinism, 1666
biology,
greenhouse strain, 303
lupin strain, 303
pollen, 303
western flower thrips, 303
BioLure,
Bactrocera, 1626
Ceratitis, 1626
nontarget, 1626
torula yeast, 1626
biomass,
dung beetles, 328
dung removal, 328
forest fragments, 328
monkey and cow dung, 328
biotic and abiotic factors,
habitat type, 57
parasitoid diversity and stem borer parasitism, 57
trophic interactions, 57
biotic interactions,
community, 798
dryland salinity, 798
environmental change, 798
temporary water, 798
biting midges,
adulticides, 591
Diptera, 591
Hymenoptera, 591
blueberry, 1428, 1990
(Z)-7-tetradecen-2-one, 1980
Anomala (=*Exomala*) *orientalis*, 1980
hand-applied plastic dispensers, 1980
harvestmen, 1428
Opiliones, 1428
species diversity, 1428
boring incitant,
bark beetle, 661
host-range prediction, 661
invasive species, 661
Pinaceae, 661
Borrelia burgdorferi,
Berberis thunbergii, 1911
Ixodes scapularis, 1911
microclimate, 1911
vapor pressure deficit, 1911

Briere-1 model,
degree-days, 1012
developmental threshold, 1012
nonlinear model, 1012
brown stink bug,
α-bisabolene epoxide, 610
green stink bug, 610
methyl (*E,E,Z*)-2,4,6-decatrienoate, 610
southern green stink bug, 610
Bt cotton, 1523, 1698
Apis mellifera, 1523
Astylus atromaculatus, 1523
Cry1Ac, 1698
gene flow, 1523
Helicoverpa armigera, 1698
insect population, 1523
resistance frequency, 1698
Bt maize, 702, 2045
Agrotis segetum, 702
Coleoptera, 2045
Cry1Ab, 702
cutworm, 702
neonicotinoid seed treatments, 2045
nontarget effects, 2045
Bt rice, 243, 715, 1369
Bacillus thuringiensis, 1369
collembolan community, 243
nontarget effect, 1369
nontarget insect, 715
nontarget organisms, 243
rice thrip, 715
risk assessment, 243
Stenchaetothrips biformis, 715
tritrophic interactions, 1369
Unmeliata insecticeps, 1369
Buchnera aphidicola,
aphidiidae, 389
development, 389
endosymbiosis, 389
life history, 389
butternut squash,
Acalymma vittatum, 1953
Cucurbita maxima, 1953
integrated pest management (IPM), 1953
pollination, 1953

C

Ca^{2+} signaling,
ethylene/jasmonic acid signaling, 1206
redox potential, 1206
Russian wheat aphid, 1206
salicylic acid signaling, 1206
Cactoblastis cactorum,
bioclimatic models, 368
cactus moth, 368
climate envelope, 368
model validation, 368
cactophilic *Drosophila*,
host shifts, 865
hybrids, 865
male genitalia, 865
wing morphology, 865
cactus moth, 368, 1884
bioclimatic models, 368
biological invasion, 1884
Cactoblastis cactorum, 368
climate envelope, 368
larval performance, 1884
model validation, 368
Opuntia, 1884
oviposition preference, 1884
Calepitrimerus vitis,
degree-days, 2006
developmental parameters, 2006
grapevine rust mite, 2006
protogyne, 2006
Campoletis chlorideae,
host discrimination, 1249
mass rearing, 1249
Mythimna separata, 1249
superparasitism, 1249
Canada,
climate change, 1122
Lepidoptera, 1122
phenology, 1122
canonical correspondence analysis,
Coccinellidae, 79
Córdoba and Granada provinces, 79
two-way indicator species analysis, 79
Cape-ivy,
German ivy, 114
risk assessment, 114
Senecio mikanioides, 114
weed biological control, 114
Carabidae,
green lacewing, 653
herbivore-induced plant volatiles, 653
HIPV, 653
lady beetles, 653
carabids,
indicator species, 738
riparian habitats, 738
saltcedar, 738
species diversity, 738
Carduus nutans, 1858, 1866
biological control, 1858
Cirsium, 1858, 1866
nontarget, 1858, 1866
Rhinocyllus conicus, 1858, 1866
Trichosirocalus horridus, 1866
Caryocar brasiliense,
ants, 1903
cerrado savanna, 1903
myrmecophily, 1903
stink bugs, 1903
cassava green mite,
Amblydromalus manihoti, 695
Euseius fustis, 695
Phytoseiidae, 695
Typhlodromalus aripo, 695
caste location,
colony composition, 295
seasonal variations, 295
termites, 295
vineyard pest, 295
cattle,
grassland, 775
grazing, 775
microclimate, 775
pest management, 775
cellulase,
bacteria, 378
gut, 378
protozoa, 378
16S rRNA gene, 378
Centaurea stoebe,
Larinus spp., 164
mowing, 164
seed production, 164
Urophora affinis, 164
cerambycid beetle,
COI, 1336
longicorn beetle, 1336
16S, 1336
wood feeder, 1336
Cerambycidae,
kairomone, 1997
pheromone, 1997
Spondylidinae, 1997
Tetropium fuscum, 1997
Ceratitis capitata,
field dispersal, 570
fruit fly, 570
sterile insect technique, 570
Ceratitis,
Bactrocera, 1626
BioLure, 1626
nontarget, 1626
torula yeast, 1626
Cerotoma trifurcata,
population dynamics, 727
seasonal abundance, 727
cerrado savanna,
ants, 1903
Caryocar brasiliense, 1903
myrmecophily, 1903
stink bugs, 1903
Ceutorhynchus obstrictus,
parasitoids, 1195
pest management, 1195
plant nutrients, 1195
spatiotemporal dynamics, 1195
chestnut weevil,
American chestnut, 787
Chinese chestnut, 787
Ormyrus labotus, 787
Torymus sinensis, 787

Chilocorus bipustulatus,
biological control, 1352
development, 1352
temperature, 1352
thermal constant, 1352
Chinese chestnut,
American chestnut, 787
chestnut weevil, 787
Ormyrus labotus, 787
Torymus sinensis, 787
Choristoneura rosaceana,
antenna, 625
Grapholita molesta, 625
pheromone, 625
sensory adaptation, 625
Chow test,
ANCOVA, 2033
hypothesis of isomorphy rate, 2033
lower developmental threshold, 2033
Chrysura kyrae,
Eastern redbud, 351
emergence models, 351
Osmia lignaria, 351
phenology, 351
Cirsium, 1858, 1866
biological control, 1858
Carduus nutans, 1858, 1866
nontarget, 1858, 1866
Rhinocyllus conicus, 1858, 1866
Trichosirocalus horridus, 1866
citrus, 618, 678
citrus pest, 678
Hemiptera, 618
Huanglongbing, 618, 678
kairomone, 618
leaf volatiles, 618
Murraya paniculata, 678
thermal requirements, 678
citrus pest,
citrus, 678
huanglongbing, 678
Murraya paniculata, 678
thermal requirements, 678
classification and regression trees,
coastline, 88
inland, 88
mosquito species, 88
Ross River virus, 88
clear-cut area,
Atlantic rainforest, 359
community structure, 359
dung beetle ecology, 359
climate,
insects, 1775
phenology, 1775
seasonality, 1775
taxa, 1775
climate change, 337, 1092, 1122, 1291
Canada, 1122
community, 337
dispersal, 1092
green leaf volatiles, 1291
habitat change, 337
herbivorous insects, 1291
insect, 337
invasive species, 1092
Lepidoptera, 1122
phenology, 1122
population, 337
relative humidity, 1092
temperature, 1092
Y-tube olfactometers, 1291
climate envelope,
bioclimatic models, 368
Cactoblastis cactorum, 368
cactus moth, 368
model validation, 368
clinal latitudinal variation,
diapause, 1751
Paralobesia viteana, 1751
viticulture, 1751
Vitis, 1751
coastline,
classification and regression trees, 88
inland, 88
mosquito species, 88
Ross River virus, 88
Coccidae,
conservation biological control, 1829
Eulecanium cerasorum, 1829
Neolecanium cornuparvum, 1829
Formicidae, 1829
Coccinella,
gut analysis, 576
Hippodamia, 576
pollen, 576
predation, 576
Coccinellidae, 79, 763
aphids, 763
biological control, 763
canonical correspondence analysis, 79
Córdoba and Granada provinces, 79
relay-intercropping, 763
sorghum, 763
two-way indicator species analysis, 79
Coccoidea,
bacteriocytes, 1178
endosymbiont, 1178
scale insects, 1178
systematics, 1178
cocoon predation,
diapause, 1893
Limacodidae, 1893
phenotypic selection analysis, 1893
pupal predation, 1893
codlemone,
apple, 1283
Cydia pomonella, 1283
mating disruption, 1283
traps, 1283
coevolution,
leaf-cutting ants, 105
parasites, 105
superinfection, 105
COI,
cerambycid beetle, 1336
longicorn beetle, 1336
S16, 1336
wood feeder, 1336
cold tolerance,
glycerol, 498
host plant effects, 498
overwintering survival, 498
supercooling point, 498
Coleoptera, 753, 2045
Bt maize, 2045
community composition, 753
Formicidae, 753
neonicotinoid seed treatments, 2045
nontarget effects, 2045
richness, 753
vegetation, 753
collembolan community,
Bt rice, 243
nontarget organisms, 243
risk assessment, 243
colony composition,
caste location, 295
seasonal variations, 295
termites, 295
vineyard pest, 295
Columbia Basin,
pheromone-baited traps, 1
Phthorimaea operculella, 1
population dynamics, 1
spatial distribution, 1
communal roosting,
Heliconius, 907
humidity, 907
light, 907
wing pattern, 907
community, 337, 798
biotic interactions, 798
climate change, 337
dryland salinity, 798
environmental change, 798
habitat change, 337
insect, 337
population, 337
temporary water, 798

community analysis,
bark beetle, 406
microbial diversity, 406
symbiosis, 406
community composition,
Coleoptera, 753
Formicidae, 753
richness, 753
vegetation, 753
community ecology,
competition, 1756
functional groups, 1756
savanna grasslands, 1756
soil ecology, 1756
community structure,
Atlantic rainforest, 359
clear-cut area, 359
dung beetle ecology, 359
compensation,
artificial injury, 528
rice, 528
sugarcane borer, 528
competition, 346, 1165, 1756
community ecology, 1756
dung, 1165
functional groups, 1756
grasshoppers, 346
herbivory, 346
lottery dynamic, 1165
nidification, 1165
Orthoptera, 346
phenology, 346
savanna grasslands, 1756
soil ecology, 1756
tunneler beetles, 1165
competitive displacement,
aggregation, 1794
Allee effect, 1794
SADIE analysis, 1794
spatial association, 1794
Conotrachelus nenuphar,
mitochondrial cytochrome oxidase I, 396
reproductive incompatibility, 396
Wolbachia, wsp, 396
conservation,
biodiversity, 1816
biological control, 1816
Elymus trachycaulus, 1816
generalist predators, 1816
IPM, 1816
western corn rootworm, 1816
conservation biocontrol,
manipulative biocontrol, 939
nectar source, 939
conservation biological control,
Coccidae, 1329
Eulecanium cerasorum,1829
Neolecanium cornuparvum, 1829
Formicidae, 1829
consumption,
Encelia, 159
herbivory, 159
coprophilous,
baited pitfall traps, 1151
beetles, 1151
forest types, 1151
necrophagous, 1151
copulatory behavior,
Amara, 1554
feeding ecology, 1554
Harpalus (*Pseduoophonus*), 1554
predispersal seed predation, 1554
corn,
host suitability, 1085
Lygus lineolaris, 1085
Zea mays, 1085
Corythuca arcuata,
lacebug, 492
light, 492
oak, 492
Tingidae, 492
Costa Rica,
biological control, 1848
host-specificity, 1848
invasive species, 1848
life history, 1848
cotton, 459, 1878
androcyclic mode, 1878
anholocyclic mode, 1878
Aphis gossypii, 1878
genetically modified viruses, 459
Helicoverpa armigera, 459
intermediate mode, 1878
mixed infection, 459
transmission, 459
cotton aphid,
ant–aphid interaction, 1492
Argentine ant, 1492
green peach aphid, 1492
odorous house ant, 1492
cover crops,
native grass, 970
natural enemies, 970
saltbush, 970
vineyards, 970
critical daylength response,
biological control, 1666
biological invasions, 1666
voltinism, 1666
crowding,
nitrogen, 685
plant quality, 685
wing dimorphism, 685
wing polymorphism, 685
Cry1Ab,
Agrotis segetum, 702
Bt maize, 702
cutworm, 702
Cry1Ac,
Bt cotton, 1698
Helicoverpa armigera, 1698
resistance frequency, 1698
Cucurbita maxima,
Acalymma vittatum, 1953
butternut squash, 1953
integrated pest management (IPM), 1953
pollination, 1953
Cucurbita pepo ssp,
Acalymma vittatum, 140
bacterial wilt disease, 140
real-time polymerase chain reaction, 140
texana, 140
Erwinia tracheiphila, 140
Culex annulirostris,
kairomones, 545
Melanotaenia duboulayi, 545
mosquito, 545
oviposition, 545
Culex,
egg retention, 190
West Nile virus, 190
Curculionidae,
exotic species, 1242
Phyllobius oblongus, 1242
Polydrusus sericeus, 1242
root herbivores, 1242
cutworm,
Agrotis segetum, 702
Bt maize, 702
Cry1Ab, 702
Cyclocephala,
Paenibacillus, 1838
Popillia japonica, 1838
Serratia, 1838
Stictospora, 1838
Cydia molesta,
Grapholita molesta, 561
microhabitat, 561
temperature, 561
thermotaxis, 561
Cydia pomonella, 583, 1283, 1399
apple, 583, 1283
areawide, 1399
codlemone, 1283
geostatistics, 1399
immunomarking, 1399
lures, 583
mark-capture, 1399
mating disruption, 1283
monitoring, 583
traps, 1283
Cynanchum, 68
domestic survey, 68
natural enemies, 68
Vincetoxicum, 68
weed biological control, 68

cytoplasmic incompatibility,
infection density, 1801
Nilaparvata lugens, 1801
quantitative polymerase chain reaction, 1801
Wolbachia, 1801
Córdoba and Granada provinces,
canonical correspondence analysis, 79
Coccinellidae, 79
two-way indicator species analysis, 79

D

defoliation,
elm spanworm, 121
Ennomos subsignarius, 121
fecundity, 121
host species, 121
degree-days, 202, 1012, 1323, 2006
alfalfa insects, 202
alfalfa management, 202
Asian longhorned beetle, 1323
Briere-1 model, 1012
Calepitrimerus vitis, 2006
development, 1323
developmental parameters, 2006
developmental threshold, 1012
grapevine rust mite, 2006
Hypera postica, 202
nonlinear model, 1012
protogyne, 2006
pupation, 1323
survival, 1323
temperature, 1323
demographic expansion,
Dendroctonus pseudotsugae, 1685
divergence time, 1685
glaciations, 1685
range expansion, 1685
demographic parameters,
development period, 1389
diel activity, 1389
oviposition period, 1389
Scirtothrips dorsalis, 1389
Dendroctonus pseudotsugae,
demographic expansion, 1685
divergence time, 1685
glaciations, 1685
range expansion, 1685
Dendroctonus,
nitrogen fertilization, 821
Ophiostoma, 821
symbiotic fungi, 821
temperature, 821
dendroentomology,
Agrilus planipennis, 253
biological invasion, 253
emerald ash borer, 253
eradication, 253
density dependence,
grasshoppers, 312
intraspecific competition, 312
metabolism, 312
temperature, 312
development, 389, 883, 1021, 1323, 1352, 1545, 1676
Ambrosia artemisiifolia, 1021
aphidiidae, 389
Asian longhorned beetle, 1323
biological control, 1352
Buchnera aphidicola, 389
Chilocorus bipustulatus, 1352
degree-days, 1323
endosymbiosis, 389
fecundity, 1021, 1676
Homalodisca vitripennis, 1545
host plant, 1545
inducible defense, 883
interspecific competition, 883
life history, 389
life table, 1545
life table parameters, 1676
Octodonta nipae, 1676
Ophraella communa, 1021
population growth, 1676
pupation, 1323
reproduction, 883, 1545
survival, 1323
systemic transportation, 883
temperature, 1021, 1323, 1352
thermal constant, 1352
development period,
demographic parameters, 1389
diel activity, 1389
oviposition period, 1389
Scirtothrips dorsalis, 1389
development time,
alfalfa weevil, 177
immature survival, 177
modeling, 177
rate isomorphy, 177
developmental parameters,
Calepitrimerus vitis, 2006
degree-days, 2006
grapevine rust mite, 2006
protogyne, 2006
developmental rate,
Aulacorthum solani, 1631
foxglove aphid, 1631
greenhouse floriculture pests, 1631
Lactin model, 1631
developmental threshold,
Briere-1 model, 1012
degree-days, 1012
nonlinear model, 1012
developmental time, 999, 2039
Anagrus nilaparvatae, 2039
fecundity, 2039
life table, 999
population, 999
Rhynchophorus ferrugineus, 999
successive generation, 2039
temperature, 999
transgenic Bt rice, 2039
Diabrotica virgifera virgifera,
belowground interactions, 1532
Fusarium verticillioides, 1532
larval development, 1532
root colonization, 1532
diapause, 1068, 1751, 1893
clinal latitudinal variation, 1751
cocoon predation, 1893
Helicoverpa armigera, 1068
Limacodidae, 1893
overwintering, 1068
Paralobesia viteana, 1751
phenotypic selection analysis, 1893
pupal predation, 1893
survival rate, 1068
traps, 1068
viticulture, 1751
Vitis, 1751
Diaprepes abbreviatus,
diet formulation, 670
mixture optimization, 670
response surface model, 670
diel activity,
demographic parameters, 1389
development period, 1389
oviposition period, 1389
Scirtothrips dorsalis, 1389
diet formulation,
Diaprepes abbreviatus, 670
mixture optimization, 670
response surface model, 670
diffusion model,
dispersal kernel, 1101
Stomoxys, 1101
dimorphic heteroecy,
host maturity, 484
leaf abscission, 484
migration, 484
soybean, 484
Diptera, 591, 835
adulticides, 591
biting midges, 591
Hymenoptera, 591
Macrotermes gilvus, 835
Misotermes mindeni, 835
parasitism, 835
termites, 835
disease risk,
immune defense, 1715
microorganism, 1715
pathogen, 1715
social insect, 1715

dispersal, 898, 1092
climate change, 1092
Hylobius warreni, 898
invasive species, 1092
mark-recapture, 898
mark-release, 898
matrix habitat, 898
relative humidity, 1092
temperature, 1092
dispersal kernel,
diffusion model, 1101
Stomoxys, 1101
dispersion,
aggregation, 1420
edge effect, 1420
spatial-temporal variation, 1420
displacement,
B biotype-specific primers, 1028
Bemisia tabaci, 1028
invasion biology, 1028
divergence time,
demographic expansion, 1685
Dendroctonus pseudotsugae, 1685
glaciations, 1685
range expansion, 1685
diversity,
environmental variables, 1744
hierarchical partitioning analysis, 1744
domestic survey,
Cynanchum, 68
natural enemies, 68
Vincetoxicum, 68
weed biological control, 68
drift behavior,
growth rate, 2017
temperature, 2017
Drosophilidae,
ammonium, 989
attractant, 989
nontarget, 989
putrescine, 989
dryland salinity,
biotic interactions, 798
community, 798
environmental change, 798
temporary water, 798
dung, 811, 1165
assemblage structure, 811
competition, 1165
Kalahari, 811
landscape, 811
lottery dynamic, 1165
nidification, 1165
Tswalu, 811
tunneler beetles, 1165
dung beetle ecology,
Atlantic rainforest, 359
clear-cut area, 359
community structure, 359
dung beetles,
biomass, 328
dung removal, 328
forest fragments, 328
monkey and cow dung, 328
dung removal,
biomass, 328
dung beetles, 328
forest fragments, 328
monkey and cow dung, 328
Dyar's rule,
European grapevine moth, 1652
larval instars, 1652
mandible length, 1652

E

early immigration,
global warming, 1705
light trap catches, 1705
Nilaparvata lugens, 1705
Eastern redbud,
Chrysura kyrae, 351
emergence models, 351
Osmia lignaria, 351
phenology, 351
ecological impact,
environmental risk assessment, 1359
genetic engineering, 1359
transgenic field trial, 1359
edge effect,
aggregation, 1420
dispersion, 1420
spatial-temporal variation, 1420
egg clutch size,
glassy-winged sharpshooter, 15
host variety, 15
seasonal phenology, 15
smoke tree sharpshooter, 15
egg retention,
Culex, 190
West Nile virus, 190
electroantennogram,
Aegorhinus superciliosus, 1276
behavioral bioassay, 1276
R-limonene, 1276
elm spanworm,
defoliation, 121
Ennomos subsignarius, 121
fecundity, 121
host species, 121
Elymus trachycaulus,
biodiversity, 1816
biological control, 1816
conservation, 1816
generalist predators, 1816
IPM, 1816
western corn rootworm, 1816
emerald ash borer,
Agrilus planipennis, 253
biological invasion, 253
dendroentomology, 253
eradication, 253
emergence,
fruit fly parasitoid, 1315
matric potential, 1315
Mediterranean fruit fly, 1315
pupal stage duration, 1315
emergence models,
Chrysura kyrae, 351
Eastern redbud, 351
Osmia lignaria, 351
phenology, 351
Encarsia,
Bemisia tabaci, 912
Eretmocerus, 912
sweetpotato whitefly, 912
Encelia,
consumption, 159
herbivory, 159
endo-brevicomin,
attractant, 633
bark beetle pheromone, 633
exo-brevicomin, 633
gas chromatography-electroantennographic detection, 633
endosymbiont,
bacteriocytes, 1178
Coccoidea, 1178
scale insects, 1178
systematics, 1178
endosymbiosis,
aphidiidae, 389
Buchnera aphidicola, 389
development, 389
life history, 389
Ennomos subsignarius,
defoliation, 121
elm spanworm, 121
fecundity, 121
host species, 121
environmental change,
biotic interactions, 798
community, 798
dryland salinity, 798
temporary water, 798
environmental limitation,
abiotic interactions, 1659
invasive species, 1659
low temperature, 1659
range limitation, 1659
environmental risk assessment,
ecological impact, 1359
genetic engineering, 1359
transgenic field trial, 1359
environmental variables,
diversity, 1744
hierarchical partitioning analysis, 1744

epidemiology,
Allium cepa, 266
management, 266
Thrips tabaci, 266
Tospovirus, 266
eradication,
Agrilus planipennis, 253
biological invasion, 253
dendroentomology, 253
emerald ash borer, 253
Eretmocerus,
Bemisia tabaci, 912
Encarsia, 912
sweetpotato whitefly, 912
Eriosoma lanigerum,
phenology, 286
woolly apple aphid, 286
Erwinia tracheiphila,
Acalymma vittatum, 140
bacterial wilt disease, 140
Cucurbita pepo ssp, 140
real-time polymerase chain reaction, 140
texana, 140
establishment,
Adelges tsugae, 440
Laricobius nigrinus, 440
release strategy, 440
Tsuga canadensis, 440
ethylene/jasmonic acid signaling,
Ca^{2+} signaling, 1206
redox potential, 1206
Russian wheat aphid, 1206
salicylic acid signaling, 1206
Eucharitidae,
agroforest, 1473
biodiversity, 1473
poneromorph, 1473
trophic interaction, 1473
Eulecanium cerasorum, 1473
Neolecanium cornuparvum, 1473
Coccidae, 1829
conservation biological control, 1829
Formicidae, 1829
Eumeninae,
acarinarium, 424
guarding mutualism, 424
host control, 424
parasite, 424
European grapevine moth,
Dyar's rule, 1652
larval instars, 1652
mandible length, 1652
Euschistus servus, 1184, 1765
Nezara viridula, 1184, 1765
planting date, 1765
Podisus maculiventris, 1184
Telenomus podisi, 1184
Trichopoda pennipes, 1765
Trissolcus basalis, 1184
Euseius fustis,
Amblydromalus manihoti, 695
cassava green mite, 695
Phytoseiidae, 695
Typhlodromalus aripo, 695
exo-brevicomin,
attractant, 633
bark beetle pheromone, 633
endo-brevicomin, 633
gas chromatography-electroantennographic detection, 633
exotic species, 1242, 1539
Acer saccaharum, 1539
Curculionidae, 1242
folivore, 1539
Phyllobius oblongus, 1242
plant-insect interactions, 1539
Polydrusus sericeus, 1242
root herbivores, 1242
Rubus, 1539

F

fecundity, 121, 195, 892, 1004, 1021, 1676, 2025, 2039
abiotic factors, 195
aggregation, 892
Ambrosia artemisiifolia, 1021
Anagrus nilaparvatae, 2039
defoliation, 121
development, 1021, 1676
developmental time, 2039
elm spanworm, 121
Ennomos subsignarius, 121
grain, 1004
host species, 121
intrinsic rate of increase, 2025
Japanese beetle, 892
life table parameters, 1676
lipid, 892
Logan-6, 2025
longevity, 1004
mate choice, 892
net reproductive rate, 2025
Octodonta nipae, 1676
Ophraella communa, 1021
oviposition, 2025
oviposition period, 1004
population growth, 1676
stored products, 1004
successive generation, 2039
temperature, 1021
transgenic Bt rice, 2039
vegetable leafminer, 195
feeding,
glassy-winged sharpshooter, 1264
mortality, 1264
overwintering, 1264
temperature, 1264
feeding ecology,
Amara, 1554
copulatory behavior, 1554
Harpalus (*Pseduoophonus*), 1554
predispersal seed predation, 1554
field dispersal,
Ceratitis capitata, 570
fruit fly, 570
sterile insect technique, 570
Fiorinia externa,
Adelges tsugae, 513
herbivory, 513
Tsuga canadensis, 513
fire,
livestock grazing, 1441
Orthoptera, 1441
population dynamics, 1441
fire ant,
phorid, 1593
Pseudacteon, 1593
Solenopsis, 1593
flight,
mark-release-recapture, 930
pheromones, 930
stored products, 930
traps, 930
flight activity,
habitat, 1482
oak–hickory forest, 1482
Xyleborus, 1482
Xylosandrus, 1482
flight potential,
larval density, 1579
Loxostege sticticalis, 1579
preoviposition period, 1579
triglyceride contents, 1579
foliar chemicals,
performance, 1990
Raphanus sativus, 1990
spatial variation, 1990
Spodoptera litura, 1990
folivore,
Acer saccaharum, 1539
exotic species, 1539
plant-insect interactions, 1539
Rubus, 1539
foraging,
immunomarking, 1936
odorous house ant, 1936
protein marking, 1936
Tapinoma sessile, 1936
trophallaxis, 1936
forecast model,
Markov chain theory, 1737
Nilaparvata lugens, 1737
rice, 1737
Sogatella furcifera, 1737
forest fragments,
biomass, 328
dung beetles, 328

dung removal, 328
monkey and cow dung, 328
forest regeneration,
Hylobius warreni, 476
Pinus contorta, 476
root form, 476
tree stability, 476
forest types,
baited pitfall traps, 1151
beetles, 1151
coprophilous, 1151
necrophagous, 1151
Formicidae, 753, 1447, 1829
Acadia National Park, 1447
Coccidae, 1829
Coleoptera, 753
community composition, 753
conservation biological control, 1829
Eulecanium cerasorum, 1829
Myrmica rubra, 1447
Neolecanium cornuparvum, 1829
richness, 753
species, 1447
survey, 1447
vegetation, 753
foxglove aphid,
Aulacorthum solani, 1631
developmental rate, 1631
greenhouse floriculture pests, 1631
Lactin model, 1631
Frankliniella fusca,
life table, 505
onion, 505
pollen, 505
thrips reproduction, 505
free amino acids,
antibiosis, 856
Aphis glycines resistance, 856
fruit fly,
Ceratitis capitata, 570
field dispersal, 570
sterile insect technique, 570
fruit fly parasitoid,
emergence, 1315
matric potential, 1315
Mediterranean fruit fly, 1315
pupal stage duration, 1315
functional groups,
community ecology, 1756
competition, 1756
savanna grasslands, 1756
soil ecology, 1756
functional response, 1570, 1586
Aphis fabae, 1586
biological control, 1570
handling time, 1586
Hippodamia variegata, 1586
searching efficiency, 1586
temperature-dependent development, 1570
Fusarium verticillioides,
belowground interactions, 1532
Diabrotica virgifera virgifera, 1532
larval development, 1532
root colonization, 1532

G

galling insects,
Amazon, 1134
biodiversity, 1134
mining, 1134
restoration ecology, 1134
gas chromatography-electroantennographic detection, 600, 633
attractant, 633
bark beetle pheromone, 633
behavioral experiments, 600
endo-brevicomin, 633
exo-brevicomin, 633
Helicoverpa armigera, 600
Microplitics mediator, 600
plant volatiles, 600
gene flow, 232, 1523
amplified fragment length polymorphism, 232
Apis mellifera, 1523
Astylus atromaculatus, 1523
Bt cotton, 1523
geographic population, 232
insect population, 1523
Loxostege sticticalis, 232
migration, 232
generalist predators,
biodiversity, 1816
biological control, 1816
conservation, 1816
Elymus trachycaulus, 1816
IPM, 1816
western corn rootworm, 1816
genetic engineering,
ecological impact, 1359
environmental risk assessment, 1359
transgenic field trial, 1359
genetic structure,
amplified fragment length polymorphism markers, 1043
insect pest, 1043
pink hibiscus mealybug, 1043
polymorphism, 1043
genetic variation,
mtDNA, 1344
Pectinophora gossypiella, 1344
population structure, 1344
genetically modified crop, 707, 1051
Bacillus thuringiensis, 707
nontarget effect, 707, 1051
phosphorus, 1051
relative food consumption rate, 1051
relative metabolic rate, 1051
stream ecology, 707
genetically modified viruses,
cotton, 459
Helicoverpa armigera, 459
mixed infection, 459
transmission, 459
geographic information system,
geostatistics, 956
SADIE, 956
sampling, 956
spatial association, 956
geographic population,
amplified fragment length polymorphism, 232
gene flow, 232
Loxostege sticticalis, 232
migration, 232
geostatistics, 944, 956, 1399, 1943
areawide, 1399
Cydia pomonella, 1399
geographic information system, 956
immunomarking, 1399
inverted distance weighted, 944, 1943
mark-capture, 1399
patchiness regression, 944, 1943
SADIE, 956
sampling, 944, 956, 1943
spatial association, 956
Taylor's Power Law, 944, 1943
Geotrupinae,
Mexico, 42
predictive distribution models, 42
species richness distribution, 42
German ivy, 114, 841
Cape-ivy, 114
host specificity, 841
nontarget impacts, 841
risk assessment, 114
Senecio mikanioides, 114, 841
weed biological control, 114, 841
glaciations,
demographic expansion, 1685
Dendroctonus pseudotsugae, 1685
divergence time, 1685
range expansion, 1685

glassy-winged sharpshooter, 15, 1264
egg clutch size, 15
feeding, 1264
host variety, 15
mortality, 1264
overwintering, 1264
seasonal phenology, 15
smoke tree sharpshooter, 15
temperature, 1264
global warming, 1643, 1705
Acrididae, 1643
early immigration, 1705
insect life cycles, 1643
light trap catches, 1705
Nilaparvata lugens, 1705
Orthoptera, 1643
voltinism, 1643
glycerol,
cold tolerance, 498
host plant effects, 498
overwintering survival, 498
supercooling point, 498
Glycine max,
Lygus lineolaris, 1111
soybeans, 1111
tarnished plant bug, 1111
grain,
fecundity, 1004
longevity, 1004
oviposition period, 1004
stored products, 1004
grape plants,
leaf temperature, 1232
plant water stress, 1232
predatory mites, 1232
spider mites, 1232
grapevine rust mite,
Calepitrimerus vitis, 2006
degree-days, 2006
developmental parameters, 2006
protogyne, 2006
Grapholita molesta, 561, 625
antenna, 625
Choristoneura rosaceana, 625
Cydia molesta, 561
microhabitat, 561
pheromone, 625
sensory adaptation, 625
temperature, 561
thermotaxis, 561
grasshoppers, 312, 346
competition, 346
density dependence, 312
herbivory, 346
intraspecific competition, 312
metabolism, 312
Orthoptera, 346
phenology, 346
temperature, 312
grassland,
cattle, 775
grazing, 775
microclimate, 775
pest management, 775
grassland spatial structure,
ant nest–seedling spatial relationship, 127
Pogonomyrmex barbatus, 127
Ripley's K function, 127
seedling establishment, 127
grazing,
cattle, 775
grassland, 775
microclimate, 775
pest management, 775
green lacewing,
Carabidae, 653
herbivore-induced plant volatiles, 653
HIPV, 653
lady beetles, 653
green leaf volatiles,
climate change, 1291
herbivorous insects, 1291
Y-tube olfactometers, 1291
green peach aphid,
ant–aphid interaction, 1492
Argentine ant, 1492
cotton aphid, 1492
odorous house ant, 1492
green stink bug,
brown stink bug, 610
methyl (*E,E,Z*)-2,4,6-decatrienoate, 610
southern green stink bug, 610
α-bisabolene epoxide, 610
greenhouse floriculture pests,
Aulacorthum solani, 1631
developmental rate, 1631
foxglove aphid, 1631
Lactin model, 1631
greenhouse strain,
biology, 303
lupin strain, 303
pollen, 303
western flower thrips, 303
growth rate,
drift behavior, 2017
temperature, 2017
guarding mutualism,
acarinarium, 424
Eumeninae, 424
host control, 424
parasite, 424
gut,
bacteria, 378
cellulase, 378
protozoa, 378
16S rRNA gene, 378
gut analysis,
Coccinella, 576
Hippodamia, 576
pollen, 576
predation, 576
gut flora accommodation,
ampicillin, 1170
oxytetracycline, 1170
tylosin, 1170
Wolbachia, 1170

H

habitat,
flight activity, 1482
oak–hickory forest, 1482
Xyleborus, 1482
Xylosandrus, 1482
habitat change,
climate change, 337
community, 337
insect, 337
population, 337
habitat characteristics of *Nannophya pygmaea*,
Juncus effusus, 278
oligotrophic state of water, 278
succession in abandoned paddy fields, 278
habitat quality,
air movement, 919
movement behavior, 919
shelter, 919
Tribolium castaneum, 919
habitat type,
biotic and abiotic factors, 57
parasitoid diversity and stem borer parasitism, 57
trophic interactions, 57
hand-applied plastic dispensers,
(Z)-7-tetradecen-2-one, 1980
Anomala (=*Exomala*) *orientalis*, 1980
blueberries, 1980
handling time,
Aphis fabae, 1586
functional response, 1586
Hippodamia variegata, 1586
searching efficiency, 1586
Harpalus (*Pseduoophonus*),
Amara, 1554
copulatory behavior, 1554
feeding ecology, 1554
predispersal seed predation, 1554
harvestmen,
blueberry, 1428
Opiliones, 1428
species diversity, 1428

Hawaiian archipelago,
biological control, 430
nontarget, 430
parasitism rates, 430
hedgerow,
Aphis glycines, 50
model selection, 50
Rhamnus cathartica, 50
Heliconius,
communal roosting, 907
humidity, 907
light, 907
wing pattern, 907
Helicoverpa armigera, 459, 552, 600, 1068, 1698
behavioral experiments, 600
behavioral trade-offs, 552
Bt cotton, 1698
cotton, 459
Cry1Ac, 1698
diapause, 1068
gas chromatography–electroantennographic detection, 600
genetically modified viruses, 459
larval performance, 552
Microplitics mediator, 600
mixed infection, 459
overwintering, 1068
oviposition preference, 552
plant volatiles, 600
realized fecundity, 552
resistance frequency, 1698
survival rate, 1068
transmission, 459
traps, 1068
Hemiptera,
citrus, 618
Huanglongbing, 618
kairomone, 618
leaf volatiles, 618
herbivore-induced plant volatiles,
Carabidae, 653
green lacewing, 653
HIPV, 653
lady beetles, 653
herbivorous insects,
climate change, 1291
green leaf volatiles, 1291
Y-tube olfactometers, 1291
herbivory, 159, 346, 513, 849
Adelges tsugae, 513
competition, 346
consumption, 159
Encelia, 159
Fiorinia externa, 513
grasshoppers, 346
insect–plant relationship, 849
leaf traits, 849
Orthoptera, 346
phenology, 346
plant quality, 849
Tsuga canadensis, 513
Heteroptera,
phenology, 1724
population ecology, 1724
seasonality, 1724
sex ratio, 1724
hierarchical partitioning analysis,
diversity, 1744
environmental variables, 1744
Hippodamia variegata,
Aphis fabae, 1586
functional response, 1586
handling time, 1586
searching efficiency, 1586
Hippodamia,
Coccinella, 576
gut analysis, 576
pollen, 576
predation, 576
HIPV,
Carabidae, 653
green lacewing, 653
herbivore-induced plant volatiles, 653
lady beetles, 653
Homalodisca vitripennis,
development, 1545
host plant, 1545
life table, 1545
reproduction, 1545
honey bee,
Aethina tumida, 1731
Kenya, 1731
Kodamaea ohmeri, 1731
Pheidole megacephala, 1731
horizontal transmission,
intestinal flora, 98
proctodeal trophallaxis, 98
symbiosis, 98
host control,
acarinarium, 424
Eumeninae, 424
guarding mutualism, 424
parasite, 424
host discrimination,
Campoletis chlorideae, 1249
mass rearing, 1249
Mythimna separata, 1249
superparasitism, 1249
host maturity,
dimorphic heteroecy, 484
leaf abscission, 484
migration, 484
soybean, 484
host orientation,
Asiatic rice borer, 1929
host range, 1929
Lepidoptera, 1929
olfaction, 1929
host plant,
development, 1545
Homalodisca vitripennis, 1545
life table, 1545
reproduction, 1545
host plant effects,
cold tolerance, 498
glycerol, 498
overwintering survival, 498
supercooling point, 498
host range,
Asiatic rice borer, 1929
host orientation, 1929
Lepidoptera, 1929
olfaction, 1929
host shifts,
cactophilic *Drosophila*, 865
hybrids, 865
male genitalia, 865
wing morphology, 865
host species,
defoliation, 121
elm spanworm, 121
Ennomos subsignarius, 121
fecundity, 121
host specificity,
nontarget impacts, 841
Senecio mikanioides, 841
German ivy, 841
weed biological control, 841
host suitability, 449, 1085
biological control, 449
corn, 1085
Lygus lineolaris, 1085
parasitoid, 449
stored-product insects, 449
wasp, 449
Zea mays, 1085
host susceptibility,
host vigor, 520
phloem thickness, 520
red oak borer, 520
host tree defense,
parasitoids, 1513
predators, 1513
Spathius agrili, 1513
Tetrastichus planipennisi, 1513
host variety,
egg clutch size, 15
glassy-winged sharpshooter, 15
seasonal phenology, 15
smoke tree sharpshooter, 15
host vigor,
host susceptibility, 520
phloem thickness, 520
red oak borer, 520
host-range prediction,
bark beetle, 661
boring incitant, 661
invasive species, 661
Pinaceae, 661

host-specificity,
biological control, 1848
Costa Rica, 1848
invasive species, 1848
life history, 1848
hosts, 1457, 1506
Anastrepha, 1457
biological control, 1506
parasitism, 1506
parasitoids, 1457, 1506
Tephritidae, 1457
Huanglongbing, 618, 678
citrus, 618, 678
citrus pest, 678
Hemiptera, 618
kairomone, 618
leaf volatiles, 618
Murraya paniculata, 678
thermal requirements, 678
humidity,
communal roosting, 907
Heliconius, 907
light, 907
wing pattern, 907
hybrids,
cactophilic *Drosophila*, 865
host shifts, 865
male genitalia, 865
wing morphology, 865
Hylobius warreni, 476, 898
dispersal, 898
forest regeneration, 476
mark-recapture, 898
mark-release, 898
matrix habitat, 898
Pinus contorta, 476
root form, 476
tree stability, 476
Hymenoptera,
adulticides, 591
biting midges, 591
Diptera, 591
Hypera postica,
alfalfa insects, 202
alfalfa management, 202
degree-day, 202
hyperspectral,
arthropod damage, 1255
multispectral, 1255
hypothesis of isomorphy rate,
ANCOVA, 2033
Chow test, 2033
lower developmental threshold, 2033

I

immature survival,
alfalfa weevil, 177
development time, 177
modeling, 177
rate isomorphy, 177
immune defense,
disease risk, 1715
microorganism, 1715
pathogen, 1715
social insect, 1715
immunomarking, 1399, 1936
areawide, 1399
Cydia pomonella, 1399
foraging, 1936
geostatistics, 1399
mark-capture, 1399
odorous house ant, 1936
protein marking, 1936
Tapinoma sessile, 1936
trophallaxis, 1936
indicator species,
carabids, 738
riparian habitats, 738
saltcedar, 738
species diversity, 738
indirect effects,
above- and below-ground interactions, 1561
induced plant responses, 1561
phytoparasite, 1561
root herbivory, 1561
induced plant responses,
above- and below-ground interactions, 1561
indirect effects, 1561
phytoparasite, 1561
root herbivory, 1561
inducible defense,
development, 883
interspecific competition, 883
reproduction, 883
systemic transportation, 883
infection density,
cytoplasmic incompatibility, 1801
Nilaparvata lugens, 1801
quantitative polymerase chain reaction, 1801
Wolbachia, 1801
inflorescences,
microhabitat use, 320
Oxyopidae, 320
sun exposure, 320
tropical dry forest, 320
inland,
classification and regression trees, 88
coastline, 88
mosquito species, 88
Ross River virus, 88
insect,
climate change, 337
community, 337
habitat change, 337
population, 337
insect extract,
mate location, 1302
olfactometer, 1302
pheromone, 1302
potato, 1302
insect life cycles,
Acrididae, 1643
global warming, 1643
Orthoptera, 1643
voltinism, 1643
insect pest,
amplified fragment length polymorphism markers, 1043
genetic structure, 1043
pink hibiscus mealybug, 1043
polymorphism, 1043
insect population,
Apis mellifera, 1523
Astylus atromaculatus, 1523
Bt cotton, 1523
gene flow, 1523
insecticide,
parasitoid, 535
sublethal effect, 535
T. evanescens, 535
T. semblidis, 535
insect–plant relationship,
herbivory, 849
insect–plant relationship, 849
leaf traits, 849
plant quality, 849
insects,
climate, 1775
phenology, 1775
seasonality, 1775
taxa, 1775
integrated pest management, 1617, 1953
absolute sampling, 1617
Acalymma vittatum, 1953
butternut squash, 1953
Cucurbita maxima, 1953
pollination, 1953
relative sampling, 1617
wing dimorphism, 1617
wing diphenism, 1617
intermediate mode,
androcyclic mode, 1878
anholocyclic mode, 1878
Aphis gossypii, 1878
cotton, 1878
interspecific competition,
development, 883
inducible defense, 883
reproduction, 883
systemic transportation, 883
intestinal flora,
horizontal transmission, 98
proctodeal trophallaxis, 98
symbiosis, 98

intraspecific competition,
density dependence, 312
grasshoppers, 312
metabolism, 312
temperature, 312
intrinsic rate of increase,
fecundity, 2025
Logan-6, 2025
net reproductive rate, 2025
oviposition, 2025
invasion biology,
B biotype-specific primers, 1028
Bemisia tabaci, 1028
displacement, 1028
invasive species, 661, 1092, 1601, 1659, 1848
abiotic interactions, 1659
agonistic behavior, 1601
bark beetle, 661
biological control, 1848
boring incitant, 661
climate change, 1092
Costa Rica, 1848
dispersal, 1092
environmental limitation, 1659
host-range prediction, 661
host-specificity, 1848
life history, 1848
low temperature, 1659
Pinaceae, 661
range limitation, 1659
relative humidity, 1092
subterranean termite, 1601
temperature, 1092
tunnel blockage, 1601
inverted distance weighted, 944, 1943
geostatistics, 944, 1943
patchiness regression, 944, 1943
sampling, 944, 1943
Taylor's Power Law, 944, 1943
IPM,
biodiversity, 1816
biological control, 1816
conservation, 1816
Elymus trachycaulus, 1816
generalist predators, 1816
western corn rootworm, 1816
Ips pini,
aggregation pheromone, 874
ipsdienol, 874
monoterpenes, 874
variance components, 874
ipsdienol,
aggregation pheromone, 874
Ips pini, 874
monoterpenes, 874
variance components, 874
isotope discrimination factor,
atmospheric nitrogen, 1810
stable isotope ratios, 1810
termites, 1810
Ixodes scapularis,
Berberis thunbergii, 1911
Borrelia burgdorferi, 1911
microclimate, 1911
vapor pressure deficit, 1911

J

Japanese beetle,
aggregation, 892
fecundity, 892
lipid, 892
mate choice, 892
Juan Fernández Archipelago,
arthropod communities, 1159
plant invasion, 1159
trophic levels, 1159
Juncus effusus,
habitat characteristics of *Nannophya pygmaea*, 278
oligotrophic state of water, 278
succession in abandoned paddy fields, 278
juvenile hormone mimic,
Metarhizium anisopliae, 979
organic farming, 979
plant-based pesticide, 979
root maggot, 979

K

kairomone, 545, 618, 642, 1973, 1997
aggregation pheromones, 642
Cerambycidae, 1997
citrus, 618
Culex annulirostris, 545
Hemiptera, 618
Huanglongbing, 618
leaf volatiles, 618
Melanotaenia duboulayi, 545
mosquito, 545
oviposition, 545
pheromone, 1997
physiological condition, 1973
Scolytidae, 642
Scolytus multistriatus, 642
Scolytus schevyrewi, 642
soybean, 1973
Spondylidinae, 1997
synthetic pheromone, 1973
Tetropium fuscum, 1997
Kalahari,
assemblage structure, 811
dung, 811
landscape, 811
Tswalu, 811
Kenya,
Aethina tumida, 1731
honey bee, 1731
Kodamaea ohmeri, 1731
Pheidole megacephala, 1731
Kodamaea ohmeri,
Aethina tumida, 1731
honey bee, 1731
Kenya, 1731
Pheidole megacephala, 1731

L

lacebug,
Corythuca arcuata, 492
light, 492
oak, 492
Tingidae, 492
Lactin model,
Aulacorthum solani, 1631
developmental rate, 1631
foxglove aphid, 1631
greenhouse floriculture pests, 1631
lady beetles,
Carabidae, 653
green lacewing, 653
herbivore-induced plant volatiles, 653
HIPV, 653
landing counts,
Asian tiger mosquito, 1608
aspirator, 1608
BG-Sentinel, 1608
surveillance, 1608
landscape,
assemblage structure, 811
dung, 811
Kalahari, 811
Tswalu, 811
landscape ecology,
Aphis glycines, 31
biological control, 31
plant chemistry, 31
potassium, 31
Laricobius nigrinus,
Adelges tsugae, 440
establishment, 440
release strategy, 440
Tsuga canadensis, 440
Larinus spp.,
Centaurea stoebe, 164
mowing, 164
seed production, 164
Urophora affinis, 164
larval competition,
Aphidius ervi, 1500
biological control, 1500
Macrosiphum euphorbiae, 1500
Praon volucre, 1500
larval density,
flight potential, 1579
Loxostege sticticalis, 1579

preoviposition period, 1579
triglyceride contents, 1579
larval development,
belowground interactions, 1532
Diabrotica virgifera virgifera, 1532
Fusarium verticillioides, 1532
root colonization, 1532
larval instars,
Dyar's rule, 1652
European grapevine moth, 1652
mandible length, 1652
larval performance, 552, 1884
behavioral trade-offs, 552
biological invasion, 1884
cactus moth, 1884
Helicoverpa armigera, 552
Opuntia, 1884
oviposition preference, 552, 1884
realized fecundity, 552
Lasius grandis,
ant activity, 827
biological control, 827
Pheidole pallidula, 827
Plagiolepis schmitzii, 827
Lasius neoniger, 1309, 1788
ants, 1788
ant-tending, 1309
aphid-tending, 1788
Aphis glycines, 1309, 1788
Scymnus louisianae, 1309
leaf abscission,
dimorphic heteroecy, 484
host maturity, 484
migration, 484
soybean, 484
leaf temperature,
grape plants, 1232
plant water stress, 1232
predatory mites, 1232
spider mites, 1232
leaf traits,
herbivory, 849
insect–plant relationship, 849
plant quality, 849
leaf volatiles,
citrus, 618
Hemiptera, 618
Huanglongbing, 618
kairomone, 618
leaf-cutting ants,
coevolution, 105
parasites, 105
superinfection, 105
leafminer,
autoinoculation device, 468
Beauveria bassiana, 468
Liriomyza huidobrensis, 468
Metarhizium anisopliae, 468
Lepidoptera, 1122, 1929
Asiatic rice borer, 1929
Canada, 1122
climate change, 1122
host orientation, 1929
host range, 1929
olfaction, 1929
phenology, 1122
Leptographium spp,
bark beetles, 415
Ophiostomatoid fungi, 415
Pinus palustris, 415
regeneration weevils, 415
life history, 389, 1848
aphidiidae, 389
biological control, 1848
Buchnera aphidicola, 389
Costa Rica, 1848
development, 389
endosymbiosis, 389
host-specificity, 1848
invasive species, 1848
life table, 505, 999, 1545
development, 1545
developmental time, 999
Frankliniella fusca, 505
Homalodisca vitripennis, 1545
host plant, 1545
onion, 505
pollen, 505
population, 999
reproduction, 1545
Rhynchophorus ferrugineus, 999
temperature, 999
thrips reproduction, 505
life table parameters,
development, 1676
fecundity, 1676
Octodonta nipae, 1676
population growth, 1676
light, 492, 907
communal roosting, 907
Corythuca arcuata, 492
Heliconius, 907
humidity, 907
lacebug, 492
oak, 492
Tingidae, 492
wing pattern, 907
light trap catches,
early immigration, 1705
global warming, 1705
Nilaparvata lugens, 1705
Limacodidae,
cocoon predation, 1893
diapause, 1893
phenotypic selection analysis, 1893
pupal predation, 1893
Linepithema humile,
arboreal foraging, 1141
Argentine ants, 1141
biological invasion, 1141
Pinus taeda, 1141
lipid,
aggregation, 892
fecundity, 892
Japanese beetle, 892
mate choice, 892
Liriomyza huidobrensis,
autoinoculation device, 468
Beauveria bassiana, 468
leafminer, 468
Metarhizium anisopliae, 468
livestock grazing,
fire, 1441
Orthoptera, 1441
population dynamics, 1441
locomotor activity,
behavior, 1922
oviposition, 1922
rotation resistance, 1922
soybean herbivory, 1922
Logan-6,
fecundity, 2025
intrinsic rate of increase, 2025
net reproductive rate, 2025
oviposition, 2025
longevity,
fecundity, 1004
grain, 1004
oviposition period, 1004
stored products, 1004
longicorn beetle,
cerambycid beetle, 1336
COI, 1336
S16, 1336
wood feeder, 1336
lottery dynamic,
competition, 1165
dung, 1165
nidification, 1165
tunneler beetles, 1165
low temperature,
abiotic interactions, 1659
environmental limitation, 1659
invasive species, 1659
range limitation, 1659
lower developmental threshold,
ANCOVA, 2033
Chow test, 2033
hypothesis of isomorphy rate, 2033
Loxostege sticticalis, 232, 1579
amplified fragment length polymorphism, 232
flight potential, 1579
gene flow, 232
geographic population, 232
larval density, 1579
migration, 232

preoviposition period, 1579
triglyceride contents, 1579
lupin strain,
biology, 303
greenhouse strain, 303
pollen, 303
western flower thrips, 303
lures,
apple, 583
Cydia pomonella, 583
monitoring, 583
Lygus lineolaris, 1085, 1111
corn, 1085
Glycine max, 1111
host suitability, 1085
soybeans, 1111
tarnished plant bug, 1111
Zea mays, 1085

M

Macrosiphum euphorbiae,
Aphidius ervi, 1500
biological control, 1500
larval competition, 1500
Praon volucre, 1500
Macrotermes gilvus,
Diptera, 835
Misotermes mindeni, 835
parasitism, 835
termites, 835
male genitalia,
cactophilic *Drosophila*, 865
host shifts, 865
hybrids, 865
wing morphology, 865
male-produced pheromone,
Anoplophora glabripennis, 169
monitoring traps, 169
plant volatiles, 169
management,
Allium cepa, 266
epidemiology, 266
Thrips tabaci, 266
Tospovirus, 266
mandible length,
Dyar's rule, 1652
European grapevine moth, 1652
larval instars, 1652
Mangifera indica,
adult emergence, 1409
population dynamics, 1409
thrips, 1409
manipulative biocontrol,
conservation biocontrol, 939
nectar source, 939
mark-capture,
areawide, 1399
Cydia pomonella, 1399
geostatistics, 1399
immunomarking, 1399
mark-recapture,
dispersal, 898
Hylobius warreni, 898
mark-release, 898
matrix habitat, 898
mark-release,
dispersal, 898
Hylobius warreni, 898
mark-recapture, 898
matrix habitat, 898
mark-release-recapture,
flight, 930
pheromones, 930
stored products, 930
traps, 930
Markov chain theory,
forecast model, 1737
Nilaparvata lugens, 1737
rice, 1737
Sogatella furcifera, 1737
mass rearing,
Campoletis chlorideae, 1249
host discrimination, 1249
Mythimna separata, 1249
superparasitism, 1249
mate choice,
aggregation, 892
fecundity, 892
Japanese beetle, 892
lipid, 892
mate location,
insect extract, 1302
olfactometer, 1302
pheromone, 1302
potato, 1302
mating disruption,
apple, 1283
codlemone, 1283
Cydia pomonella, 1283
traps, 1283
matric potential,
emergence, 1315
fruit fly parasitoid, 1315
Mediterranean fruit fly, 1315
pupal stage duration, 1315
matrix habitat,
dispersal, 898
Hylobius warreni, 898
mark-recapture, 898
mark-release, 898
Medicago sativa,
alfalfa leafcutting bee, 149
bee density, 149
reproductive success, 149
Mediterranean fruit fly,
emergence, 1315
fruit fly parasitoid, 1315
matric potential, 1315
pupal stage duration, 1315
Melanotaenia duboulayi,
Culex annulirostris, 545
kairomones, 545
mosquito, 545
oviposition, 545
metabolism,
density dependence, 312
grasshoppers, 312
intraspecific competition, 312
temperature, 312
Metarhizium anisopliae, organic farming,
juvenile hormone mimic, 979
plant-based pesticide, 979
root maggot, 979
Metarhizium anisopliae,
autoinoculation device, 468
Beauveria bassiana, 468
leafminer, 468
Liriomyza huidobrensis, 468
methyl (*E,E,Z*)-2,4,6-decatrienoate,
brown stink bug, 610
green stink bug, 610
methyl (*E,E,Z*)-2,4,6-decatrienoate, 610
southern green stink bug, 610
α-bisabolene epoxide, 610
Mexico,
Geotrupinae, 42
predictive distribution models, 42
species richness distribution, 42
Miami blue butterfly,
aerial ULV application, 1961
mosquito control, 1961
naled, 1961
nontarget impact, 1961
microbial diversity,
bark beetle, 406
community analysis, 406
symbiosis, 406
microclimate, 775, 1911
Berberis thunbergii, 1911
Borrelia burgdorferi, 1911
cattle, 775
grassland, 775
grazing, 775
Ixodes scapularis, 1911
pest management, 775
vapor pressure deficit, 1911
microhabitat,
Cydia molesta, 561
Grapholita molesta, 561
temperature, 561
thermotaxis, 561
microhabitat use,
inflorescences, 320
Oxyopidae, 320
sun exposure, 320
tropical dry forest, 320
microorganism,
disease risk, 1715
immune defense, 1715

pathogen, 1715
social insect, 1715
Microplitics mediator,
behavioral experiments, 600
gas chromatography–electroantennographic detection, 600
Helicoverpa armigera, 600
plant volatiles, 600
midge mouth parts,
river pollution, 210
water and sediments quality, 210
migration, 232, 484
amplified fragment length polymorphism, 232
dimorphic heteroecy, 484
gene flow, 232
geographic population, 232
host maturity, 484
leaf abscission, 484
Loxostege sticticalis, 232
soybean, 484
mining,
Amazon, 1134
biodiversity, 1134
galling insects, 1134
restoration ecology, 1134
Misotermes mindeni,
Diptera, 835
Macrotermes gilvus, 835
parasitism, 835
termites, 835
mitochondrial cytochrome oxidase I,
Conotrachelus nenuphar, 396
reproductive incompatibility, 396
Wolbachia, wsp, 396
mixed infection,
cotton, 459
genetically modified viruses, 459
Helicoverpa armigera, 459
transmission, 459
mixture optimization,
Diaprepes abbreviatus, 670
diet formulation, 670
response surface model, 670
model selection,
Aphis glycines, 50
hedgerow, 50
Rhamnus cathartica, 50
model validation,
bioclimatic models, 368
Cactoblastis cactorum, 368
cactus moth, 368
climate envelope, 368
modeling,
alfalfa weevil, 177
development time, 177
immature survival, 177
rate isomorphy, 177
monitoring,
apple, 583
Cydia pomonella, 583
lures, 583
monitoring traps,
Anoplophora glabripennis, 169
male-produced pheromone, 169
plant volatiles, 169
monkey and cow dung,
biomass, 328
dung beetles, 328
dung removal, 328
forest fragments, 328
monoterpenes,
aggregation pheromone, 874
Ips pini, 874
ipsdienol, 874
variance components, 874
mortality,
feeding, 1264
glassy-winged sharpshooter, 1264
overwintering, 1264
temperature, 1264
mosquito,
Culex annulirostris, 545
kairomones, 545
Melanotaenia duboulayi, 545
oviposition, 545
mosquito control,
aerial ULV application, 1961
Miami blue butterfly, 1961
naled, 1961
nontarget impact, 1961
mosquito species,
classification and regression trees, 88
coastline, 88
inland, 88
Ross River virus, 88
movement behavior,
air movement, 919
habitat quality, 919
shelter, 919
Tribolium castaneum, 919
mowing,
Centaurea stoebe, 164
Larinus spp., 164
seed production, 164
Urophora affinis, 164
mtDNA,
genetic variation, 1344
Pectinophora gossypiella, 1344
population structure, 1344
multispectral,
arthropod damage, 1255
hyperspectral, 1255
Murraya paniculata,
citrus, 678
citrus pest, 678
huanglongbing, 678
thermal requirements, 678
myrmecophily,
ants, 1903
Caryocar brasiliense, 1903
cerrado savanna, 1903
stink bugs, 1903
Myrmica rubra,
Acadia National Park, 1447
Formicidae, 1447
species, 1447
survey, 1447
Mythimna separata,
Campoletis chlorideae, 1249
host discrimination, 1249
mass rearing, 1249
superparasitism, 1249

N

naled,
aerial ULV application, 1961
Miami blue butterfly, 1961
mosquito control, 1961
nontarget impact, 1961
native grass,
cover crops, 970
natural enemies, 970
saltbush, 970
vineyards, 970
natural enemies, 68, 970, 1466
cover crops, 970
domestic survey, 68
native grass, 970
predator-prey interactions, 1466
saltbush, 970
trophic interactions, 1466
Vincetoxicum Cynanchum, 68
vineyards, 970
web location, 1466
web-weaving spiders, 1466
weed biological control, 68
navel orangeworm,
almonds, 1059
overwintering emergence, 1059
pistachios, 1059
sex ratio, 1059
necrophagous,
baited pitfall traps, 1151
beetles, 1151
coprophilous, 1151
forest types, 1151
nectar source,
conservation biocontrol, 939
manipulative biocontrol, 939
neonicotinoid seed treatments,
Bt maize, 2045
Coleoptera, 2045
nontarget effects, 2045
net reproductive rate,
fecundity, 2025
intrinsic rate of increase, 2025

Logan-6, 2025
oviposition, 2025
Nezara viridula, 1184, 1765
Euschistus servus, 1184, 1765
planting date, 1765
Podisus maculiventris, 1184
Telenomus podisi, 1184
Trichopoda pennipes, 1765
Trissolcus basalis, 1184
nidification,
competition, 1165
dung, 1165
lottery dynamic, 1165
tunneler beetles, 1165
Nilaparvata lugens, 1705, 1737, 1801
cytoplasmic incompatibility, 1801
early immigration, 1705
forecast model, 1737
global warming, 1705
infection density, 1801
light trap catches, 1705
Markov chain theory, 1737
quantitative polymerase chain reaction, 1801
rice, 1737
Sogatella furcifera, 1737
Wolbachia, 1801
nitrogen,
crowding, 685
plant quality, 685
wing dimorphism, 685
wing polymorphism, 685
nitrogen fertilization,
Dendroctonus, 821
Ophiostoma, 821
symbiotic fungi, 821
temperature, 821
nonlinear model,
Briere-1 model, 1012
degree-days, 1012
developmental threshold, 1012
nontarget, 430, 989, 1626, 1858, 1866
ammonium, 989
attractant, 989
Bactrocera, 1626
biological control, 430, 1858
BioLure, 1626
Carduus nutans, 1858, 1866
Ceratitis, 1626
Cirsium, 1858, 1866
Drosophilidae, 989
Hawaiian archipelago, 430
parasitism rates, 430
putrescine, 989
Rhinocyllus conicus, 1858, 1866
torula yeast, 1626
Trichosirocalus horridus, 1866
nontarget effect, 707, 1051, 1369, 2045
Bacillus thuringiensis, 707, 1369
Bt maize, 2045
Bt rice, 1369
Coleoptera, 2045
genetically modified crop, 707, 1051
neonicotinoid seed treatments, 2045
relative food consumption rate, 1051
relative metabolic rate, 1051
stream ecology, 707
tritrophic interactions, 1369
Ummeliata insecticeps, 1369
nontarget impact, 841, 1961
aerial ULV application, 1961
German ivy, 841
host specificity, 841
Miami blue butterfly, 1961
mosquito control, 1961
naled, 1961
Senecio mikanioides, 841
weed biological control, 841
nontarget insect,
Bt rice, 715
rice thrip, 715
Stenchaetothrips biformis, 715
nontarget organisms,
Bt rice, 243
collembolan community, 243
risk assessment, 243
nontarget pest,
Adelphocoris suturalis Jakovlev, 1378
oviposition behavior, 1378
population density, 1378
transgenic cotton, 1378

O

oak,
Corythuca arcuata, 492
lacebug, 492
light, 492
Tingidae, 492
oak–hickory forest,
flight activity, 1482
habitat, 1482
Xyleborus, 1482
Xylosandrus, 1482
Octodonta nipae,
development, 1676
fecundity, 1676
life table parameters, 1676
population growth, 1676
odorous house ant, 1492, 1936
ant–aphid interaction, 1492
Argentine ant, 1492
cotton aphid, 1492
foraging, 1936
green peach aphid, 1492
immunomarking, 1936
protein marking, 1936
Tapinoma sessile, 1936
trophallaxis, 1936
olfaction,
Asiatic rice borer, 1929
host orientation, 1929
host range, 1929
Lepidoptera, 1929
olfactometer,
insect extract, 1302
mate location, 1302
pheromone, 1302
potato, 1302
oligotrophic state of water,
habitat characteristics of *Nannophya pygmaea*, 278
Juncus effusus, 278
succession in abandoned paddy fields, 278
onion,
Frankliniella fusca, 505
life table, 505
pollen, 505
thrips reproduction, 505
Ophiostoma,
Dendroctonus, 821
nitrogen fertilization, 821
symbiotic fungi, 821
temperature, 821
Ophiostomatoid fungi,
bark beetles, 415
Leptographium spp, 415
Pinus palustris, 415
regeneration weevils, 415
Ophraella communa,
Ambrosia artemisiifolia, 1021
development, 1021
fecundity, 1021
temperature, 1021
Opiliones,
blueberry, 1428
harvestmen, 1428
species diversity, 1428
Opuntia,
biological invasion, 1884
cactus moth, 1884
larval performance, 1884
oviposition preference, 1884
organic farming
juvenile hormone mimic, 979
Metarhizium anisopliae, 979
root maggot, 979
plant-based pesticide, 979
Ormyrus labotus,
American chestnut, 787
chestnut weevil, 787
Chinese chestnut, 787
Torymus sinensis, 787

Orthoptera, 346, 1441, 1643
Acrididae, 1643
competition, 346
fire, 1441
global warming, 1643
grasshoppers, 346
herbivory, 346
insect life cycles, 1643
livestock grazing, 1441
phenology, 346
population dynamics, 1441
voltinism, 1643
Osmia lignaria,
Chrysura kyrae, 351
Eastern redbud, 351
emergence models, 351
phenology, 351
overwintering, 1068, 1264
diapause, 1068
feeding, 1264
glassy-winged sharpshooter, 1264
Helicoverpa armigera, 1068
mortality, 1264
survival rate, 1068
temperature, 1264
traps, 1068
overwintering emergence,
almonds, 1059
navel orangeworm, 1059
pistachios, 1059
sex ratio, 1059
overwintering survival,
cold tolerance, 498
glycerol, 498
host plant effects, 498
supercooling point, 498
oviposition, 545, 1922, 2025
behavior, 1922
Culex annulirostris, 545
fecundity, 2025
intrinsic rate of increase, 2025
kairomones, 545
locomotor activity, 1922
Logan-6, 2025
Melanotaenia duboulayi, 545
mosquito, 545
net reproductive rate, 2025
rotation resistance, 1922
soybean herbivory, 1922
oviposition behavior,
Adelphocoris suturalis Jakovlev, 1378
nontarget pest, 1378
population density, 1378
transgenic cotton, 1378
oviposition period, 1004, 1389
demographic parameters, 1389
development period, 1389
diel activity, 1389
fecundity, 1004
grain, 1004
longevity, 1004
Scirtothrips dorsalis, 1389
stored products, 1004
oviposition preference, 552, 1884
behavioral trade-offs, 552
biological invasion, 1884
cactus moth, 1884
Helicoverpa armigera, 552
larval performance, 552, 1884
Opuntia, 1884
realized fecundity, 552
Oxyopidae,
inflorescences, 320
microhabitat use, 320
sun exposure, 320
tropical dry forest, 320
oxytetracycline,
ampicillin, 1170
gut flora accommodation, 1170
tylosin, 1170
Wolbachia, 1170

P

Paenibacillus,
Cyclocephala, 1838
Popillia japonica, 1838
Serratia, 1838
Stictospora, 1838
Paralobesia viteana,
clinal latitudinal variation, 1751
diapause, 1751
viticulture, 1751
Vitis, 1751
parasite, 105, 424
acarinarium, 424
coevolution, 105
Eumeninae, 424
guarding mutualism, 424
host control, 424
leaf-cutting ants, 105
superinfection, 105
parasitism, 835, 1506
biological control, 1506
Diptera, 835
hosts, 1506
Macrotermes gilvus, 835
Misotermes mindeni, 835
parasitoids, 1506
termites, 835
parasitism rates,
biological control, 430
Hawaiian archipelago, 430
nontarget, 430
parasitoid diversity and stem borer parasitism,
biotic and abiotic factors, 57
habitat type, 57
trophic interactions, 57
parasitoid, 449, 553, 1195, 1457, 1506, 1513
Anastrepha, 1457
biological control, 449, 1506
Ceutorhynchus obstrictus, 1195
host tree defense, 1513
hosts, 1457, 1506
host suitability, 449
insecticide, 535
parasitism, 1506
pest management, 1195
plant nutrients, 1195
predators, 1513
Spathius agrili, 1513
spatiotemporal dynamics, 1195
stored-product insects, 449
sublethal effect, 535
T. evanescens, 535
T. semblidis, 535
Tephritidae, 1457
Tetrastichus planipennisi, 1513
wasp, 449
patchiness regression, 944, 1943
geostatistics, 944, 1943
inverted distance weighted, 944, 1943
sampling, 944, 1943
Taylor's Power Law, 944, 1943
pathogen,
disease risk, 1715
immune defense, 1715
microorganism, 1715
social insect, 1715
Pectinophora gossypiella,
genetic variation, 1344
mtDNA, 1344
population structure, 1344
performance,
foliar chemicals, 1990
Raphanus sativus, 1990
spatial variation, 1990
Spodoptera litura, 1990
pest management, 775, 1195
cattle, 775
Ceutorhynchus obstrictus, 1195
grassland, 775
grazing, 775
microclimate, 775
parasitoids, 1195
plant nutrients, 1195
spatiotemporal dynamics, 1195
Pheidole megacephala,
Aethina tumida, 1731
honey bee, 1731
Kenya, 1731
Kodamaea ohmeri, 1731
Pheidole pallidula,
ant activity, 827
biological control, 827
Lasius grandis, 827
Plagiolepis schmitzii, 827

phenology, 286, 346, 351, 1122, 1724, 1775
Canada, 1122
Chrysura kyrae, 351
climate, 1775
climate change, 1122
competition, 346
Eastern redbud, 351
emergence models, 351
Eriosoma lanigerum, 286
grasshoppers, 346
herbivory, 346
Heteroptera, 1724
insects, 1775
Lepidoptera, 1122
Orthoptera, 346
Osmia lignaria, 351
population ecology, 1724
seasonality, 1724, 1775
sex ratio, 1724
taxa, 1775
woolly apple aphid, 286
phenotypic selection analysis,
cocoon predation, 1893
diapause, 1893
Limacodidae, 1893
pupal predation, 1893
pheromone, 625, 930, 1302, 1997
antenna, 625
Cerambycidae, 1997
Choristoneura rosaceana, 625
flight, 930
Grapholita molesta, 625
insect extract, 1302
kairomone, 1997
mark-release-recapture, 930
mate location, 1302
olfactometer, 1302
potato, 1302
sensory adaptation, 625
Spondylidinae, 1997
stored products, 930
Tetropium fuscum, 1997
traps, 930
pheromone-baited traps,
Columbia Basin, 1
Phthorimaea operculella, 1
population dynamics, 1
spatial distribution, 1
phloem thickness,
host susceptibility, 520
host vigor, 520
red oak borer, 520
phorid,
fire ant, 1593
Pseudacteon, 1593
Solenopsis, 1593
phosphorus,
genetically modified crop, 1051
nontarget effect, 1051
relative food consumption rate, 1051
relative metabolic rate, 1051
Phthorimaea operculella,
Columbia Basin, 1
pheromone-baited traps, 1
population dynamics, 1
spatial distribution, 1
Phyllobius oblongus,
Curculionidae, 1242
exotic species, 1242
Polydrusus sericeus, 1242
root herbivores, 1242
phylogeography,
amplified fragment length polymorphisms, 1037
potato, 1037
potato tubermoth, 1037
physiological condition,
kairomone, 1973
soybean, 1973
synthetic pheromone, 1973
phytoparasite,
above- and below-ground interactions, 1561
indirect effects, 1561
induced plant responses, 1561
root herbivory, 1561
Phytoseiidae,
Amblydromalus manihoti, 695
cassava green mite, 695
Euseius fustis, 695
Typhlodromalus aripo, 695
Pinaceae,
bark beetle, 661
boring incitant, 661
host-range prediction, 661
invasive species, 661
pink hibiscus mealybug,
amplified fragment length polymorphism markers, 1043
genetic structure, 1043
insect pest, 1043
polymorphism, 1043
Pinus contorta,
forest regeneration, 476
Hylobius warreni, 476
root form, 476
tree stability, 476
Pinus palustris,
bark beetles, 415
Leptographium spp, 415
Ophiostomatoid fungi, 415
regeneration weevils, 415
Pinus taeda,
arboreal foraging, 1141
Argentine ants, 1141
biological invasion, 1141
Linepithema humile, 1141
pistachios,
almonds, 1059
navel orangeworm, 1059
overwintering emergence, 1059
sex ratio, 1059
Plagiolepis schmitzii,
ant activity, 827
biological control, 827
Lasius grandis, 827
Pheidole pallidula, 827
plant chemistry,
Aphis glycines, 31
biological control, 31
landscape ecology, 31
potassium, 31
plant invasion,
arthropod communities, 1159
Juan Fernández Archipelago, 1159
trophic levels, 1159
plant nutrients,
Ceutorhynchus obstrictus, 1195
parasitoids, 1195
pest management, 1195
spatiotemporal dynamics, 1195
plant quality, 685, 849
crowding, 685
herbivory, 849
insect–plant relationship, 849
leaf traits, 849
nitrogen, 685
wing dimorphism, 685
wing polymorphism, 685
plant volatiles, 169, 600
Anoplophora glabripennis, 169
behavioral experiments, 600
gas chromatography–electroantennographic detection, 600
Helicoverpa armigera, 600
male-produced pheromone, 169
Microplitics mediator, 600
monitoring traps, 169
plant water stress,
grape plants, 1232
leaf temperature, 1232
predatory mites, 1232
spider mites, 1232
plant-based pesticide,
juvenile hormone mimic, 979
Metarhizium anisopliae, 979
organic farming, 979
root maggot, 979
plant-insect interactions,
Acer saccharum, 1539
exotic species, 1539
folivore, 1539
Rubus, 1539

planting date,
Euschistus servus, 1765
Nezara viridula, 1765
Trichopoda pennipes, 1765
Podisus maculiventris,
Euschistus servus, 1184
Nezara viridula, 1184
Telenomus podisi, 1184
Trissolcus basalis, 1184
Pogonomyrmex barbatus,
ant nest-seedling spatial relationship, 127
grassland spatial structure, 127
Ripley's K function, 127
seedling establishment, 127
pollen, 303, 505, 576
biology, 303
Coccinella, 576
Frankliniella fusca, 505
greenhouse strain, 303
gut analysis, 576
Hippodamia, 576
life table, 505
lupin strain, 303
onion, 505
predation, 576
thrips reproduction, 505
western flower thrips, 303
pollination,
Acalymma vittatum, 1953
butternut squash, 1953
Cucurbita maxima, 1953
integrated pest management (IPM), 1953
Polydrusus sericeus,
Curculionidae, 1242
exotic species, 1242
Phyllobius oblongus, 1242
root herbivores, 1242
polymorphism,
amplified fragment length polymorphism markers, 1043
genetic structure, 1043
insect pest, 1043
pink hibiscus mealybug, 1043
poneromorph,
agroforest, 1473
biodiversity, 1473
Eucharitidae, 1473
trophic interaction, 1473
Popillia japonica,
Cyclocephala, 1838
Paenibacillus, 1838
Serratia, 1838
Stictospora, 1838
population, 337, 999
climate change, 337
community, 337
developmental time, 999
habitat change, 337
insect, 337
life table, 999
Rhynchophorus ferrugineus, 999
temperature, 999
population density,
Adelphocoris suturalis Jakovlev, 1378
nontarget pest, 1378
oviposition behavior, 1378
transgenic cotton, 1378
population dynamics, 1, 727, 1409, 1441
adult emergence, 1409
Cerotoma trifurcata, 727
Columbia Basin, 1
fire, 1441
livestock grazing, 1441
Mangifera indica, 1409
Orthoptera, 1441
pheromone-baited traps, 1
Phthorimaea operculella, 1
seasonal abundance, 727
spatial distribution, 1
thrips, 1409
population ecology,
Heteroptera, 1724
phenology, 1724
seasonality, 1724
sex ratio, 1724
population growth,
development, 1676
fecundity, 1676
life table parameters, 1676
Octodonta nipae, 1676
population structure,
genetic variation, 1344
mtDNA, 1344
Pectinophora gossypiella, 1344
potassium,
Aphis glycines, 31
biological control, 31
landscape ecology, 31
plant chemistry, 31
potato, 1037, 1302
amplified fragment length polymorphisms, 1037
insect extract, 1302
mate location, 1302
olfactometer, 1302
pheromone, 1302
phylogeography, 1037
potato tubermoth, 1037
potato tubermoth,
amplified fragment length polymorphisms, 1037
phylogeography, 1037
potato, 1037
Praon volucre,
Aphidius ervi, 1500
biological control, 1500
larval competition, 1500
Macrosiphum euphorbiae, 1500
predation,
Coccinella, 576
gut analysis, 576
Hippodamia, 576
pollen, 576
predator-prey interactions,
natural enemies, 1466
trophic interactions, 1466
web location, 1466
web-weaving spiders, 1466
predators,
host tree defense, 1513
parasitoids, 1513
Spathius agrili, 1513
Tetrastichus planipennisi, 1513
predatory mites,
grape plants, 1232
leaf temperature, 1232
plant water stress, 1232
spider mites, 1232
predictive distribution models,
Geotrupinae, 42
Mexico, 42
species richness distribution, 42
predispersal seed predation,
Amara, 1554
copulatory behavior, 1554
feeding ecology, 1554
Harpalus (*Pseduoophonus*), 1554
preoviposition period,
flight potential, 1579
larval density, 1579
Loxostege sticticalis, 1579
triglyceride contents, 1579
proctodeal trophallaxis,
horizontal transmission, 98
intestinal flora, 98
symbiosis, 98
protein marking,
foraging, 1936
immunomarking, 1936
odorous house ant, 1936
Tapinoma sessile, 1936
trophallaxis, 1936
protogyne,
Calepitrimerus vitis, 2006
degree-days, 2006
developmental parameters, 2006
grapevine rust mite, 2006
protozoa,
bacteria, 378
cellulase, 378
gut, 378
16S rRNA gene, 378
Pseudacteon,
fire ant, 1593
phorid, 1593
Solenopsis, 1593

pupal predation,
cocoon predation, 1893
diapause, 1893
Limacodidae, 1893
phenotypic selection analysis, 1893
pupal stage duration,
emergence, 1315
fruit fly parasitoid, 1315
matric potential, 1315
Mediterranean fruit fly, 1315
pupation,
Asian longhorned beetle, 1323
degree-days, 1323
development, 1323
survival, 1323
temperature, 1323
putrescine,
ammonium, 989
attractant, 989
Drosophilidae, 989
nontarget, 989

Q

quantitative polymerase chain reaction,
cytoplasmic incompatibility, 1801
infection density, 1801
Nilaparvata lugens, 1801
Wolbachia, 1801

R

R-limonene,
Aegorhinus superciliosus, 1276
behavioral bioassay, 1276
electroantennogram, 1276
range expansion,
demographic expansion, 1685
Dendroctonus pseudotsugae, 1685
divergence time, 1685
glaciations, 1685
range limitation,
abiotic interactions, 1659
environmental limitation, 1659
invasive species, 1659
low temperature, 1659
Raphanus sativus,
foliar chemicals, 1990
performance, 1990
spatial variation, 1990
Spodoptera litura, 1990
rate isomorphy,
alfalfa weevil, 177
development time, 177
immature survival, 177
modeling, 177
real-time polymerase chain reaction,
Acalymma vittatum, 140
bacterial wilt disease, 140
Cucurbita pepo ssp, 140
texana, 140
Erwinia tracheiphila, 140
realized fecundity,
behavioral trade-offs, 552
Helicoverpa armigera, 552
larval performance, 552
oviposition preference, 552
red oak borer,
host susceptibility, 520
host vigor, 520
phloem thickness, 520
redox potential,
Ca^{2+} signaling, 1206
ethylene/jasmonic acid signaling, 1206
Russian wheat aphid, 1206
salicylic acid signaling, 1206
regeneration weevils,
bark beetles, 415
Leptographium spp, 415
Ophiostomatoid fungi, 415
Pinus palustris, 415
relative food consumption rate,
genetically modified crop, 1051
nontarget effect, 1051
relative metabolic rate, 1051
relative humidity,
climate change, 1092
dispersal, 1092
invasive species, 1092
temperature, 1092
relative metabolic rate,
genetically modified crop, 1051
nontarget effect, 1051
relative food consumption rate, 1051
relative sampling,
absolute sampling, 1617
integrated pest management, 1617
wing dimorphism, 1617
wing diphenism, 1617
relay-intercropping,
aphids, 763
biological control, 763
Coccinellidae, 763
sorghum, 763
release strategy,
Adelges tsugae, 440
establishment, 440
Laricobius nigrinus, 440
Tsuga canadensis, 440
reproduction, 883, 1545
development, 883, 1545
Homalodisca vitripennis, 1545
host plant, 1545
inducible defense, 883
interspecific competition, 883
life table, 1545
systemic transportation, 883
reproductive incompatibility,
Conotrachelus nenuphar, 396
mitochondrial cytochrome oxidase I, 396
Wolbachia, wsp, 396
reproductive success,
alfalfa leafcutting bee, 149
bee density, 149
Medicago sativa, 149
resistance frequency,
Bt cotton, 1698
Cry1Ac, 1698
Helicoverpa armigera, 1698
response surface model,
Diaprepes abbreviatus, 670
diet formulation, 670
mixture optimization, 670
restoration ecology,
Amazon, 1134
biodiversity, 1134
galling insects, 1134
mining, 1134
Rhamnus cathartica,
Aphis glycines, 50
hedgerow, 50
model selection, 50
Rhinocyllus conicus, 1858, 1866
biological control, 1858
Carduus nutans, 1858, 1866
Cirsium, 1858, 1866
nontarget, 1858, 1866
Trichosirocalus horridus, 1866
Rhynchophorus ferrugineus,
developmental time, 999
life table, 999
population, 999
temperature, 999
rice,
artificial injury, 528
compensation, 528
forecast model, 1737
Markov chain theory, 1737
Nilaparvata lugens, 1737
Sogatella furcifera, 1737
sugarcane borer, 528
rice thrip,
Bt rice, 715
nontarget insect, 715
Stenchaetothrips biformis, 715
richness,
Coleoptera, 753
community composition, 753
Formicidae, 753
vegetation, 753
riparian habitats,
carabids, 738
indicator species, 738
saltcedar, 738
species diversity, 738

Ripley's K function,
ant nest–seedling spatial relationship, 127
grassland spatial structure, 127
Pogonomyrmex barbatus, 127
seedling establishment, 127
risk assessment, 114, 243
Bt rice, 243
Cape-ivy, 114
collembolan community, 243
German ivy, 114
nontarget organisms, 243
Senecio mikanioides, 114
weed biological control, 114
river pollution,
midge mouth parts, 210
water and sediments quality, 210
root colonization,
belowground interactions, 1532
Diabrotica virgifera virgifera, 1532
Fusarium verticillioides, 1532
larval development, 1532
root form,
forest regeneration, 476
Hylobius warreni, 476
Pinus contorta, 476
tree stability, 476
root herbivores,
Curculionidae, 1242
exotic species, 1242
Phyllobius oblongus, 1242
Polydrusus sericeus, 1242
root herbivory,
above- and below-ground interactions, 1561
indirect effects, 1561
induced plant responses, 1561
phytoparasite, 1561
root maggot,
juvenile hormone mimic, 979
Metarhizium anisopliae, 979
organic farming, 979
plant-based pesticide, 979
Ross River virus,
classification and regression trees, 88
coastline, 88
inland, 88
mosquito species, 88
rotation resistance,
behavior, 1922
locomotor activity, 1922
oviposition, 1922
soybean herbivory, 1922
16S rRNA gene,
bacteria, 378
cellulase, 378
gut, 378
protozoa, 378
Rubus,
Acer saccharum, 1539
exotic species, 1539
folivore, 1539
plant-insect interactions, 1539
Russian wheat aphid, 223, 1206
aphid diets, 223
aphid saliva collection, 223
Ca^{2+} signaling, 1206
ethylene/jasmonic acid signaling, 1206
redox potential, 1206
salicylic acid signaling, 1206
saliva proteins, 223
Triticum aestivum, 223

S

S16,
cerambycid beetle, 1336
COI, 1336
longicorn beetle, 1336
wood feeder, 1336
SADIE,
geographic information system, 956
geostatistics, 956
sampling, 956
spatial association, 956
SADIE analysis,
aggregation, 1794
Allee effect, 1794
competitive displacement, 1794
spatial association, 1794
salicylic acid signaling,
Ca^{2+} signaling, 1206
ethylene/jasmonic acid signaling, 1206
redox potential, 1206
Russian wheat aphid, 1206
saliva proteins,
aphid diets, 223
aphid saliva collection, 223
Russian wheat aphid, 223
Triticum aestivum, 223
saltbush,
cover crops, 970
native grass, 970
natural enemies, 970
vineyards, 970
saltcedar,
carabids, 738
indicator species, 738
riparian habitats, 738
species diversity, 738
sampling, 944, 956, 1943
geographic information system, 956
geostatistics, 944, 956, 1943
inverted distance weighted, 944, 1943
patchiness regression, 944, 1943
SADIE, 956
spatial association, 956
Taylor's Power Law, 944, 1943
savanna grasslands,
community ecology, 1756
competition, 1756
functional groups, 1756
soil ecology, 1756
scale insects,
bacteriocytes, 1178
Coccoidea, 1178
endosymbiont, 1178
systematics, 1178
Scirtothrips dorsalis,
demographic parameters, 1389
development period, 1389
diel activity, 1389
oviposition period, 1389
Scolytidae,
aggregation pheromones, 642
kairomones, 642
Scolytus multistriatus, 642
Scolytus schevyrewi, 642
Scolytus multistriatus,
aggregation pheromones, 642
kairomones, 642
Scolytidae, 642
Scolytus schevyrewi, 642
Scolytus schevyrewi,
aggregation pheromones, 642
kairomones, 642
Scolytidae, 642
Scolytus multistriatus, 642
Scymnus louisianae,
ant-tending, 1309
Aphis glycines, 1309
Lasius neoniger, 1309
searching efficiency,
Aphis fabae, 1586
functional response, 1586
handling time, 1586
Hippodamia variegata, 1586
seasonal abundance,
Cerotoma trifurcata, 727
population dynamics, 727
seasonal phenology,
egg clutch size, 15
glassy-winged sharpshooter, 15
host variety, 15
smoke tree sharpshooter, 15
seasonal variations,
caste location, 295
colony composition, 295
termites, 295
vineyard pest, 295
seasonality, 1724, 1775
climate, 1775
Heteroptera, 1724
insects, 1775
phenology, 1724, 1775

population ecology, 1724
sex ratio, 1724
taxa, 1775
seed production,
Centaurea stoebe, 164
Larinus spp., 164
mowing, 164
Urophora affinis, 164
seedling establishment,
ant nest–seedling spatial relationship, 127
grassland spatial structure, 127
Pogonomyrmex barbatus, 127
Ripley's K function, 127
Senecio mikanioides, 114, 841
Cape-ivy, 114
German ivy, 114, 841
host specificity, 841
nontarget impacts, 841
risk assessment, 114
weed biological control, 114, 841
sensory adaptation,
antenna, 625
Choristoneura rosaceana, 625
Grapholita molesta, 625
pheromone, 625
Serratia,
Cyclocephala, 1838
Paenibacillus, 1838
Popillia japonica, 1838
Stictospora, 1838
sex ratio, 1059, 1724
almonds, 1059
Heteroptera, 1724
navel orangeworm, 1059
overwintering emergence, 1059
phenology, 1724
pistachios, 1059
population ecology, 1724
seasonality, 1724
shelter,
air movement, 919
habitat quality, 919
movement behavior, 919
Tribolium castaneum, 919
smoke tree sharpshooter,
egg clutch size, 15
glassy-winged sharpshooter, 15
host variety, 15
seasonal phenology, 15
social insect,
disease risk, 1715
immune defense, 1715
microorganism, 1715
pathogen, 1715
Sogatella furcifera,
forecast model, 1737
Markov chain theory, 1737
Nilaparvata lugens, 1737
rice, 1737
soil ecology,
community ecology, 1756
competition, 1756
functional groups, 1756
savanna grasslands, 1756
Solenopsis,
fire ant, 1593
phorid, 1593
Pseudacteon, 1593
sorghum,
aphids, 763
biological control, 763
Coccinellidae, 763
relay-intercropping, 763
southern green stink bug,
brown stink bug, 610
green stink bug, 610
methyl (*E,E,Z*)-2,4,6-decatrienoate, 610
α-bisabolene epoxide, 610
soybean, 484, 1111, 1973
dimorphic heteroecy, 484
Glycine max, 1111
host maturity, 484
kairomone, 1973
leaf abscission, 484
Lygus lineolaris, 1111
migration, 484
physiological condition, 1973
synthetic pheromone, 1973
tarnished plant bug, 1111
soybean herbivory,
behavior, 1922
locomotor activity, 1922
oviposition, 1922
rotation resistance, 1922
Spathius agrili,
host tree defense, 1513
parasitoids, 1513
predators, 1513
Tetrastichus planipennisi, 1513
spatial association,
geographic information system, 956
geostatistics, 956
SADIE, 956
sampling, 956
spatial distribution,
Columbia Basin, 1
pheromone-baited traps, 1
Phthorimaea operculella, 1
population dynamics, 1
spatial variation,
foliar chemicals, 1990
performance, 1990
Raphanus sativus, 1990
Spodoptera litura, 1990
spatial-temporal variation,
aggregation, 1420
dispersion, 1420
edge effect, 1420
spatial association,
aggregation, 1794
Allee effect, 1794
competitive displacement, 1794
SADIE analysis, 1794
spatiotemporal dynamics,
Ceutorhynchus obstrictus, 1195
parasitoids, 1195
pest management, 1195
plant nutrients, 1195
species,
Acadia National Park, 1447
Formicidae, 1447
Myrmica rubra, 1447
survey, 1447
species diversity, 738, 1428
blueberry, 1428
carabids, 738
harvestmen, 1428
indicator species, 738
Opiliones, 1428
riparian habitats, 738
saltcedar, 738
species richness distribution,
Geotrupinae, 42
Mexico, 42
predictive distribution models, 42
spider mites,
grape plants, 1232
leaf temperature, 1232
plant water stress, 1232
predatory mites, 1232
Spodoptera litura,
foliar chemicals, 1990
performance, 1990
Raphanus sativus, 1990
spatial variation, 1990
Spondylidinae,
Cerambycidae, 1997
kairomone, 1997
pheromone, 1997
Tetropium fuscum, 1997
stable isotope ratios,
atmospheric nitrogen, 1810
isotope discrimination factor, 1810
termites, 1810
Stenchaetothrips biformis,
Bt rice, 715
nontarget insect, 715
rice thrip, 715
sterile insect technique,
Ceratitis capitata, 570
field dispersal, 570
fruit fly, 570
Stictospora,
Cyclocephala, 1838
Paenibacillus, 1838

Popillia japonica, 1838
Serratia, 1838
stink bug, 134, 1903
ants, 1903
apple, 134
Caryocar brasiliense, 1903
cerrado savanna, 1903
myrmecophily, 1903
UV radiation, 134
wound response, 134
Stomoxys,
diffusion model, 1101
dispersal kernel, 1101
stored products, 930, 1004
fecundity, 1004
flight, 930
grain, 1004
longevity, 1004
mark-release-recapture, 930
oviposition period, 1004
pheromones, 930
traps, 930
stored-product insects,
biological control, 449
host suitability, 449
parasitoid, 449
wasp, 449
stream ecology,
Bacillus thuringiensis, 707
genetically modified crops, 707
nontarget effects, 707
sublethal effect,
insecticide, 535
parasitoid, 535
T. evanescens, 535
T. semblidis, 535
subterranean termite,
agonistic behavior, 1601
invasive species, 1601
tunnel blockage, 1601
succession in abandoned paddy fields,
habitat characteristics of *Nannophya pygmaea*, 278
Juncus effusus, 278
oligotrophic state of water, 278
successive generation,
Anagrus nilaparvatae, 2039
developmental time, 2039
fecundity, 2039
transgenic Bt rice, 2039
sugarcane borer,
artificial injury, 528
compensation, 528
rice, 528
sun exposure,
inflorescences, 320
microhabitat use, 320
Oxyopidae, 320
tropical dry forest, 320
supercooling point,
cold tolerance, 498
glycerol, 498
host plant effects, 498
overwintering survival, 498
superinfection,
coevolution, 105
leaf-cutting ants, 105
parasites, 105
superparasitism,
Campoletis chlorideae, 1249
host discrimination, 1249
mass rearing, 1249
Mythimna separata, 1249
surveillance,
Asian tiger mosquito, 1608
aspirator, 1608
BG-Sentinel, 1608
landing counts, 1608
survey,
Acadia National Park, 1447
Formicidae, 1447
Myrmica rubra, 1447
species, 1447
survival,
Asian longhorned beetle, 1323
degree-days, 1323
development, 1323
pupation, 1323
temperature, 1323
survival rate,
diapause, 1068
Helicoverpa armigera, 1068
overwintering, 1068
traps, 1068
sweetpotato whitefly,
Bemisia tabaci, 912
Encarsia, 912
Eretmocerus, 912
symbiosis, 98, 406
bark beetle, 406
community analysis, 406
horizontal transmission, 98
intestinal flora, 98
microbial diversity, 406
proctodeal trophallaxis, 98
symbiotic fungi,
Dendroctonus, 821
nitrogen fertilization, 821
Ophiostoma, 821
temperature, 821
synthetic pheromone,
kairomone, 1973
physiological condition, 1973
soybean, 1973
systematics,
bacteriocytes, 1178
Coccoidea, 1178
endosymbiont, 1178
scale insects, 1178
systemic transportation,
development, 883
inducible defense, 883
interspecific competition, 883
reproduction, 883

T

T. evanescens,
insecticide, 535
parasitoid, 535
sublethal effect, 535
T. semblidis, 535
T. semblidis,
insecticide, 535
parasitoid, 535
sublethal effect, 535
T. evanescens, 535
Tapinoma sessile,
foraging, 1936
immunomarking, 1936
odorous house ant, 1936
protein marking, 1936
trophallaxis, 1936
tarnished plant bug,
Glycine max, 1111
Lygus lineolaris, 1111
soybeans, 1111
taxa,
climate, 1775
insects, 1775
phenology, 1775
seasonality, 1775
Taylor's Power Law, 944, 1943
geostatistics, 944, 1943
inverted distance weighted, 944, 1943
patchiness regression, 944, 1943
sampling, 944, 1943
Telenomus podisi,
Euschistus servus, 1184
Nezara viridula, 1184
Podisus maculiventris, 1184
Trissolcus basalis, 1184
temperature, 312, 561, 821, 999, 1021, 1092, 1264, 1323, 1352, 2017
Ambrosia artemisiifolia, 1021
Asian longhorned beetle, 1323
biological control, 1352
Chilocorus bipustulatus, 1352
climate change, 1092
Cydia molesta, 561
degree-days, 1323
Dendroctonus, 821
density dependence, 312
development, 1021, 1323, 1352
developmental time, 999
dispersal, 1092
drift behavior, 2017
fecundity, 1021
feeding, 1264

glassy-winged sharpshooter, 1264
Grapholita molesta, 561
grasshoppers, 312
growth rate, 2017
intraspecific competition, 312
invasive species, 1092
life table, 999
metabolism, 312
microhabitat, 561
mortality, 1264
nitrogen fertilization, 821
Ophiostoma, 821
Ophraella communa, 1021
overwintering, 1264
population, 999
pupation, 1323
relative humidity, 1092
Rhynchophorus ferrugineus, 999
survival, 1323
symbiotic fungi, 821
thermal constant, 1352
thermotaxis, 561
temperature-dependent development,
biological control, 1570
functional response, 1570
temporary water,
biotic interactions, 798
community, 798
dryland salinity, 798
environmental change, 798
Tephritidae,
Anastrepha, 1457
hosts, 1457
parasitoids, 1457
termites, 295, 835, 1810
atmospheric nitrogen, 1810
caste location, 295
colony composition, 295
Diptera, 835
isotope discrimination factor, 1810
Macrotermes gilvus, 835
Misotermes mindeni, 835
parasitism, 835
seasonal variations, 295
stable isotope ratios, 1810
vineyard pest, 295
Tetrastichus planipennisi,
host tree defense, 1513
parasitoids, 1513
predators, 1513
Spathius agrili, 1513
Tetropium fuscum,
Cerambycidae, 1997
kairomone, 1997
pheromone, 1997
Spondylidinae, 1997
texana,
Acalymma vittatum, 140
bacterial wilt disease, 140
Cucurbita pepo ssp, 140
Erwinia tracheiphila, 140
real-time polymerase chain reaction, 140
thermal constant,
biological control, 1352
Chilocorus bipustulatus, 1352
development, 1352
temperature, 1352
thermal requirements,
citrus, 678
citrus pest, 678
huanglongbing, 678
Murraya paniculata, 678
thermotaxis,
Cydia molesta, 561
Grapholita molesta, 561
microhabitat, 561
temperature, 561
thrips,
adult emergence, 1409
Mangifera indica, 1409
population dynamics, 1409
thrips reproduction,
Frankliniella fusca, 505
life table, 505
onion, 505
pollen, 505
Thrips tabaci,
Allium cepa, 266
epidemiology, 266
management, 266
Tospovirus, 266
Tingidae,
Corythuca arcuata, 492
lacebug, 492
light, 492
oak, 492
torula yeast,
Bactrocera, 1626
BioLure, 1626
Ceratitis, 1626
nontarget, 1626
Torymus sinensis,
American chestnut, 787
chestnut weevil, 787
Chinese chestnut, 787
Ormyrus labotus, 787
Tospovirus,
Allium cepa, 266
epidemiology, 266
management, 266
Thrips tabaci, 266
transgenic Bt rice,
Anagrus nilaparvatae, 2039
developmental time, 2039
fecundity, 2039
successive generation, 2039
transgenic cotton,
Adelphocoris suturalis Jakovlev, 1378
nontarget pest, 1378
oviposition behavior, 1378
population density, 1378
transgenic field trial,
ecological impact, 1359
environmental risk assessment, 1359
genetic engineering, 1359
transmission,
cotton, 459
genetically modified viruses, 459
Helicoverpa armigera, 459
mixed infection, 459
traps, 930, 1068, 1283
apple, 1283
codlemone, 1283
Cydia pomonella, 1283
diapause, 1068
flight, 930
Helicoverpa armigera, 1068
mark-release-recapture, 930
mating disruption, 1283
overwintering, 1068
pheromones, 930
stored products, 930
survival rate, 1068
tree stability,
forest regeneration, 476
Hylobius warreni, 476
Pinus contorta, 476
root form, 476
Tribolium castaneum,
air movement, 919
habitat quality, 919
movement behavior, 919
shelter, 919
Trichopoda pennipes,
Euschistus servus, 1765
Nezara viridula, 1765
planting date, 1765
Trichosirocalus horridus,
Carduus nutans, 1866
Cirsium, 1866
nontarget, 1866
Rhinocyllus conicus, 1866
triglyceride contents,
flight potential, 1579
larval density, 1579
Loxostege sticticalis, 1579
preoviposition period, 1579
Trissolcus basalis,
Euschistus servus, 1184
Nezara viridula, 1184
Podisus maculiventris, 1184
Telenomus podisi, 1184
Triticum aestivum,
aphid diets, 223
aphid saliva collection, 223

Russian wheat aphid, 223
saliva proteins, 223
tritrophic interactions,
Bacillus thuringiensis, 1369
Bt rice, 1369
nontarget effect, 1369
Ummeliata insecticeps, 1369
trophallaxis,
foraging, 1936
immunomarking, 1936
odorous house ant, 1936
protein marking, 1936
Tapinoma sessile, 1936
trophic interaction, 57, 1466, 1473
agroforest, 1473
biodiversity, 1473
biotic and abiotic factors, 57
Eucharitidae, 1473
habitat type, 57
natural enemies, 1466
parasitoid diversity and stem borer parasitism, 57
poneromorph, 1473
predator-prey interactions, 1466
web location, 1466
web-weaving spiders, 1466
trophic levels,
arthropod communities, 1159
Juan Fernández Archipelago, 1159
plant invasion, 1159
tropical dry forest,
inflorescences, 320
microhabitat use, 320
Oxyopidae, 320
sun exposure, 320
Tsuga canadensis, 440, 513
Adelges tsugae, 440, 513
establishment, 440
Fiorinia externa, 513
herbivory, 513
Laricobius nigrinus, 440
release strategy, 440
Tswalu,
assemblage structure, 811
dung, 811
Kalahari, 811
landscape, 811
tunnel blockage,
agonistic behavior, 1601
invasive species, 1601
subterranean termite, 1601
tunneler beetles,
competition, 1165
dung, 1165
lottery dynamic, 1165
nidification, 1165
two-way indicator species analysis,
canonical correspondence analysis, 79
Coccinellidae, 79
Córdoba and Granada provinces, 79
tylosin,
ampicillin, 1170
gut flora accommodation, 1170
oxytetracycline, 1170
Wolbachia, 1170
Typhlodromalus aripo,
Amblydromalus manihoti, 695
cassava green mite, 695
Euseius fustis, 695
Phytoseiidae, 695

U

Ummeliata insecticeps,
Bacillus thuringiensis, 1369
Bt rice, 1369
nontarget effect, 1369
tritrophic interactions, 1369
Urophora affinis,
Centaurea stoebe, 164
Larinus spp., 164
mowing, 164
seed production, 164
UV radiation,
apple, 134
stink bug, 134
wound response, 134

V

vapor pressure deficit,
Berberis thunbergii, 1911
Borrelia burgdorferi, 1911
Ixodes scapularis, 1911
microclimate, 1911
variance components,
aggregation pheromone, 874
Ips pini, 874
ipsdienol, 874
monoterpenes, 874
vegetable leafminer,
abiotic factors, 195
fecundity, 195
vegetation,
Coleoptera, 753
community composition, 753
Formicidae, richness, 753
Vincetoxicum Cynanchum,
domestic survey, 68
natural enemies, 68
weed biological control, 68
vineyard pest,
caste location, 295
colony composition, 295
seasonal variations, 295
termites, 295
vineyards,
cover crops, 970
native grass, 970
natural enemies, 970
saltbush, 970
viticulture,
clinal latitudinal variation, 1751
diapause, 1751
Paralobesia viteana, 1751
Vitis, 1751
Vitis,
clinal latitudinal variation, 1751
diapause, 1751
Paralobesia viteana, 1751
viticulture, 1751
voltinism, 1643, 1666
Acrididae, 1643
biological control, 1666
biological invasions, 1666
critical daylength response, 1666
global warming, 1643
insect life cycles, 1643
Orthoptera, 1643

W

wasp,
biological control, 449
host suitability, 449
parasitoid, 449
stored-product insects, 449
water and sediments quality,
midge mouth parts, 210
river pollution, 210
web location,
natural enemies, 1466
predator-prey interactions, 1466
trophic interactions, 1466
web-weaving spiders, 1466
web-weaving spiders,
natural enemies, 1466
predator-prey interactions, 1466
trophic interactions, 1466
web location, 1466
weed biological control, 68, 114, 841
Cape-ivy, 114
domestic survey, 68
German ivy, 114
host specificity, 841
natural enemies, 68
nontarget impacts, 841
risk assessment, 114
Senecio mikanioides, 841
German ivy, 841
Senecio mikanioides, 114
Vincetoxicum Cynanchum, 68

West Nile virus,
Culex, 190
egg retention, 190
western corn rootworm,
biodiversity, 1816
biological control, 1816
conservation, 1816
Elymus trachycaulus, 1816
generalist predators, 1816
IPM, 1816
western flower thrips,
biology, 303
greenhouse strain, 303
lupin strain, 303
pollen, 303
wing dimorphism, 685, 1617
absolute sampling, 1617
crowding, 685
integrated pest management, 1617
nitrogen, 685
plant quality, 685
relative sampling, 1617
wing diphenism, 1617
wing polymorphism, 685
wing diphenism,
absolute sampling, 1617
integrated pest management, 1617
relative sampling, 1617
wing dimorphism, 1617
wing morphology,
cactophilic *Drosophila*, 865
host shifts, 865
hybrids, 865
male genitalia, 865
wing pattern,
communal roosting, 907
Heliconius, 907
humidity, 907
light, 907
wing polymorphism,
crowding, 685
nitrogen, 685
plant quality, 685
wing dimorphism, 685
Wolbachia, wsp,
Conotrachelus nenuphar, 396
mitochondrial cytochrome oxidase I, 396
reproductive incompatibility, 396
Wolbachia, 1170, 1801
ampicillin, 1170
cytoplasmic incompatibility, 1801
gut flora accommodation, 1170
infection density, 1801
Nilaparvata lugens, 1801
oxytetracycline, 1170
quantitative polymerase chain reaction, 1801
tylosin, 1170
wood feeder,
cerambycid beetle, 1336
COI, 1336
longicorn beetle, 1336
S16, 1336
woolly apple aphid,
Eriosoma lanigerum, 286
phenology, 286
wound response,
apple, 134
stink bug, 134
UV radiation, 134

X

Xyleborus,
flight activity, 1482
habitat, 1482
oak–hickory forest, 1482
Xylosandrus, 1482
Xylosandrus,
flight activity, 1482
habitat, 1482
oak–hickory forest, 1482
Xyleborus, 1482

Y

Y-tube olfactometers,
climate change, 1291
green leaf volatiles, 1291
herbivorous insects, 1291

Z

Zea mays,
corn, 1085
host suitability, 1085
Lygus lineolaris, 1085